

(12) INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

Rec'd PCT/PTO 14 JAN 2005

(19) World Intellectual Property
Organization
International Bureau




(43) International Publication Date
22 January 2004 (22.01.2004)

PCT

10/521313
(10) International Publication Number
WO 2004/007734 A1

- (51) International Patent Classification⁷: C12N 15/85 Chang-Yuil [KR/KR]; 104-502 Daewoo Hyoryung Apt, Bangbae-dong, Seocho-gu, Seoul 137-060 (KR).
- (21) International Application Number: PCT/KR2003/001400 (74) Agents: JANG, Seongku et al.; 19th Fl., KEC Building, 275-7, Yangjae-dong, Seocho-ku, Seoul 137-130 (KR).
- (22) International Filing Date: 15 July 2003 (15.07.2003) (81) Designated States (*national*): AE, AG, AL, AM, AT, AU, AZ, BA, BB, BG, BR, BY, BZ, CA, CH, CN, CO, CR, CU, CZ, DE, DK, DM, DZ, EC, EE, ES, FI, GB, GD, GE, GH, GM, HR, HU, ID, IL, IN, IS, JP, KE, KG, KP, KZ, LC, LK, LR, LS, LT, LU, LV, MA, MD, MG, MK, MN, MW, MX, MZ, NI, NO, NZ, OM, PG, PH, PL, PT, RO, RU, SC, SD, SE, SG, SK, SL, SY, TJ, TM, TN, TR, TT, TZ, UA, UG, US, UZ, VC, VN, YU, ZA, ZM, ZW.
- (25) Filing Language: English
- (26) Publication Language: English
- (30) Priority Data:
10-2002-0041764 16 July 2002 (16.07.2002) KR
10-2003-0038012 12 June 2003 (12.06.2003) KR
- (71) Applicant (*for all designated States except US*): PANGE-NOMICS CO., LTD. [KR/KR]; #304 Biotechnology Incubating Center, Seoul National University, San 56-1, Shillim-dong, Kwanak-gu, Seoul 151-742 (KR).
- (72) Inventors; and
- (75) Inventors/Applicants (*for US only*): LEE, Joon Youb [KR/KR]; 76-503 Hyundai Apt., Apkujung-dong, Kangnam-gu, Seoul 135-110 (KR). KIM, Dong-Hyeon [KR/KR]; 1267-329, Mia-6-dong, Kangbuk-gu, Seoul 142-106 (KR). CHUNG, Yeonseok [KR/KR]; Hanyang Apt. 10-303, Apkujung-dong, Kangnam-gu, Seoul 135-110 (KR). CHANG, Sun-Young [KR/KR]; 5 Fl., #602-15, Sindaebang-dong, Dongjak-gu, Seoul 156-010 (KR). LEE, Kyung-Chul [KR/KR]; Banpo Apt. 111-203, Banpobon-dong, Seocho-gu, Seoul 137-049 (KR). KANG,
- (84) Designated States (*regional*): ARIPO patent (GH, GM, KE, LS, MW, MZ, SD, SL, SZ, TZ, UG, ZM, ZW), Eurasian patent (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM), European patent (AT, BE, BG, CH, CY, CZ, DE, DK, EE, ES, FI, FR, GB, GR, HU, IE, IT, LU, MC, NL, PT, RO, SE, SI, SK, TR), OAPI patent (BF, BJ, CF, CG, CI, CM, GA, GN, GQ, GW, ML, MR, NE, SN, TD, TG).
- Published:
— with international search report
— before the expiration of the time limit for amending the claims and to be republished in the event of receipt of amendments
- For two-letter codes and other abbreviations, refer to the "Guidance Notes on Codes and Abbreviations" appearing at the beginning of each regular issue of the PCT Gazette.

(54) Title: HER-2/NEU DNA VACCINE HAVING ANTI-CANCER ACTIVITY



(57) Abstract: The present invention relates to human Her-2/neu expressing plasmid constructs having anti-cancer activity and a DNA vaccine comprising same for preventing and/or treating cancer. The Her-2/neu DNA vaccines of the present invention can be effectively used as a therapeutic vaccine in reducing metastasis after tumor surgery or as a prophylactic vaccine for people with genetic high risk.

WO 2004/007734 A1

Her-2/neu DNA VACCINE HAVING ANTI-CANCER ACTIVITY

FIELD OF THE INVENTION

5 The present invention relates to human Her-2/neu expressing plasmid constructs having anti-cancer activity and a DNA vaccine comprising same for preventing and treating cancer.

BACKGROUND OF THE INVENTION

10 The Her-2/neu or *erbB-2* gene encodes a transmembrane protein that is a member of the type I family of growth factor receptors (Akiyama, T. et al., *Science* 232: 1644-1646, 1986). Amplification of this gene results in overexpression of the encoded 185 kDa receptor tyrosine kinase.

15 The Her-2/neu protein has been found to be amplified and overexpressed in several types of human adenocarcinomas, especially in tumors of the breast and the ovary. The overexpression was correlated with short relapse time and poor survival rate of breast cancer patients (Slamon, D. J. et al., *Science* 235: 177-182, 1987), suggesting that Her-2/neu overexpression likely plays a critical role in the development
20 of human cancers. Several lines of evidence also support a direct role of Her-2/neu in the pathogenesis and clinical aggressiveness of Her-2/neu-expressing tumors (Kobayashi H. et al., *Cancer Res.* 60: 5228-5236, 2000). For example, Herceptin, a humanized anti-Her-2/neu monoclonal antibody used for treatment of Her-2/neu-expressing tumors, has been demonstrated to bring clinical benefits in advanced breast
25 cancer patients (Ewer, M. S. et. al., *Semin. Oncol.* 26: 96, 1999). In addition, Her-2/neu-specific antibodies and T cells are detected in breast and ovarian cancer patients. Therefore, Her-2/neu oncogene is an excellent target for the development of therapeutic vaccines specific for Her-2/neu-overexpressing human cancers.

30 Since human Her-2/neu gene has tyrosine kinas activity in the intracellular domain and its overexpression itself stimulates abnormal cell division, there are several attempts to eliminate possible oncogenecity of Her-2/neu by introducing a mutation into the cytoplasmic kinase domain to inhibit tyrosine kinase activity or by constructing truncated Her-2/neu plasmids lacking the intracellular or extracellular

domain (Wei, W. I. et al., *Int. J. Cancer* 81: 748-754, 1999)

Naked plasmids are attractive candidate vectors for the development of cancer vaccines encoding tumor-associated antigens. They are relatively simple to generate and safe to administer. Because they are not proteins nor associated with a viral coat, naked nucleic acids are not generally subject to neutralizing antibody reactions that can hamper the clinical efficacy of vaccines (Hellstrom, I. and Hellstrom, K. E., *J. Immunother.* 21: 119-126, 1998). In preclinical tumor models, DNA vaccines encoding rat (Chen, Y. et al., *Cancer Res.* 58: 1965-1971, 1998) or human Her-2/neu (Pilon, S. A. et al., *J. Immunol.* 167: 3201-3206, 2001) induced preventive efficacy against Her-2/neu expressing tumor cells.

Although successful preventive efficacy against Her-2/neu expressing tumor by DNA vaccination was achieved by many earlier experiments, no successful therapeutic efficacy was reported using only Her-2/neu expressing plasmids. The difficulty lies on the slow gain of antitumor immunity due to the lag time before antigenic expression of Her-2/neu expressing plasmids, while mammary tumor grows relatively fast. Therefore, some of the Her-2/neu therapeutic vaccine experiments were conducted based on the combination of DNA and cytokine-secreting tumor cells (Chen, S. A. et al., *Clin. Cancer Res.* 6: 4381-4388, 2000), or dendritic cell (Chen, Y., *Gene Ther.* 8: 316-323, 2001).

Since a DNA vaccine has many advantages including mass-productivity, safety, and convenience (Gurunathan, S. et al., *Annu. Rev. Immunol.* 18:927-974, 2001), the present inventors have endeavored to develop Her-2/neu expressing plasmid constructs having high anti-cancer activity which can be effectively used as a DNA vaccine for preventing and treating cancer.

SUMMARY OF THE INVENTION

Accordingly, it is an object of the present invention to provide a human Her-2/neu expressing plasmid construct having high antitumor activity.

Another object of the present invention is to provide a DNA vaccine composition for preventing and/or treating cancer, comprising said plasmid construct and a pharmaceutically acceptable carrier.

An additional object of the present invention is to provide a method for

preventing and/or treating cancer, comprising the step of administering an effective amount of said DNA vaccine.

BRIEF DESCRIPTION OF THE DRAWINGS

5

The above and other objects and features of the present invention will become apparent from the following description of the invention, when taken in conjunction with the accompanying drawings, which respectively show:

10

Figs. 1a and 1b: schematic procedure for preparing recombinant human pNeu plasmid constructs and a preset immunization schedule, respectively;

ECD: extracellular domain,

TM: transmembrane domain,

ICD: intracellular domain

15

Fig. 2: representative FACS histograms of medium fluorescence intensity in each of the groups of mice vaccinated with pTV2 (A), pNeu_{TM} (B), pNeu_{ECD} (C), pNeu_{TM-gDs} (D) and pNeu_{ECD-gDs} (E), respectively;

uncolored FACS histogram: control antibody,

colored FACS histogram: anti-Her-2/neu antibody

20

Fig. 3: confocal microscopic analysis of anti-Her-2/neu antibody in mouse sera immunized with pTV2 (A), pNeu_{TM} (B) and pNeu_{ECD-gDs} (C), respectively;

Fig. 4: ⁵¹Cr-release assays for comparing cytotoxic T lymphocytes (CTL) responses induced by vaccination with pTV2 (A), pNeu_{TM} (B), pNeu_{ECD} (C), pNeu_{TM-gDs} (D) and pNeu_{ECD-gDs} (E), respectively;

25

Fig. 5: preventive antitumor immunity induced by vaccination with pNeu constructs;

A: tumor size in animal model subcutaneously injected with Her2-CT26 cells

B: survival rate in animal model intravenously injected with Her2-CT26 cells

Fig. 6: comparison of preventive antitumor immunity induced by vaccination

30

with pNeu_{ECD} and pNeu_{ECD-gDs};

A: tumor size in animal model subcutaneously injected of Her2-CT26 cells

B: survival rate in animal model intravenously injected of Her2-CT26 cells

Fig. 7: therapeutic efficacy induced by vaccination with pNeu_{ECD} or pNeu_{ECD-gDs}.

gDs;

A: 1×10^5 Her2-CT26 cells, B: 5×10^5 Her2-CT26 cells

Fig. 8: representative FACS histograms of medium fluorescence intensity in each group of mice vaccinated with PBS (A), pNeu_{ECD} (B), pNeu_{TM} (C), pCK_{ECD} (D) and pCK_{TM} (E), respectively;

Fig. 9: ⁵¹Cr-release assays for comparing CTL responses induced by vaccination with PBS (A), pNeu_{ECD} (B), pNeu_{TM} (C), pCK_{ECD} (D) and pCK_{TM} (E), respectively;

Fig. 10: preventive antitumor immunity induced by vaccination with pCK_{ECD} and pCK_{TM};

A: tumor size in animal model subcutaneously injected of Her2-CT26 cells

B: survival rate in animal model intravenously injected of Her2-CT26 cells

Fig. 11: therapeutic efficacies induced by vaccination with pCK_{ECD} and pCK_{TM};

Figs. 12a and 12b: vaccination schedule of co-injection with pCK_{TM} and cytokine plasmids and ⁵¹Cr-release assays for comparing CTL responses induced thereby, respectively;

Figs. 13a to 13d: preventive antitumor effect induced by co-injection with pCK_{TM} and cytokine plasmids, wherein 13a shows vaccination schedule; 13b, tumor size in subcutaneous injection model of Her2-CT26 cells (a parenthesis means the percentage of mouse having no tumor growth per treatment group); 13c and 13d, survival rate in intravenous injection model of Her2-CT26 cells (a parenthesis means the percentage of live mouse per treatment group);

Figs. 14a to 14c: therapeutic efficacies induced by co-injection with pCK_{TM} and pCK-cytokine plasmids, wherein 14a shows vaccination schedule; 14b and 14c, survival rate in intravenous injection model vaccination with pCK_{TM}-cytokine plasmids. of Her2-CT26 cells (a parenthesis means the percentage of live mouse per treatment group);

Figs. 15a to 15d: preventive antitumor effect induced by pCK_{TM}-cytokine plasmids, wherein 15a shows schematic procedure of constructing bicistronic plasmids; 15b, vaccination schedule; 15c, tumor size in subcutaneous injection model of Her2-CT26 cells (a parenthesis means the percentage of mouse having no tumor growth per treatment group); 15d, survival rate in intravenous injection model of Her2-CT26 cells (a parenthesis means the percentage of live mouse per treatment group);

Fig. 16: therapeutic efficacies induced by

DETAILED DESCRIPTION OF THE INVENTION

5 In accordance with one aspect of the present invention, there is provided Her-2/neu expressing plasmid constructs having anti-cancer activity which is prepared by inserting a truncated human Her-2/neu gene into pTV2 or pCK vector.

First, the present invention provides Her-2/neu expressing plasmid constructs encoding a truncated Her-2/neu gene that lacked the cytoplasmic kinase domain (intercellular domain), the truncated gene being selected because a plasmid encoding the
10 full-length human Her-2/neu may adversely affect the physiology of the cells that takes up plasmid DNA. The truncated Her-2/neu gene has the nucleotide sequence of SEQ ID NO: 2 comprising the Her-2/neu transmembrane and extracellular domains, and is inserted into pTV2 vector which gives a high expression level of a foreign gene (Lee, S. W. et al., *J. Virol.* 72: 8430-8436, 1998)

15 The present invention also provides Her-2/neu expressing plasmid constructs encoding the truncated human Her-2/neu gene of SEQ ID NO: 3 that lacks the transmembrane domain of the Her-2/neu gene of SEQ ID NO: 2, which results in the secretion of the expressed protein into the cell exterior.

Further, the present invention provides Her-2/neu expressing plasmid constructs
20 of which the signal peptide sequence is replaced by the herpes simplex virus type I glycoprotein D signal (gDs) sequence which is known to facilitate the efficient expression and secretion of human immunodeficiency virus type I gp160 (Berman, P. W. et al., *J. Virol.* 63: 3489-3498, 1989).

In a preferred embodiment of the present invention, four Her-2/neu expressing
25 plasmid constructs based pTV2 vector (pNeu_{TM}, pNeu_{ECD}, pNeu_{TM-gDs} and pNeu_{ECD-gDs}) are generated encoding either the Her-2/neu transmembrane and extracellular domains (pNeu_{TM} and pNeu_{TM-gDs}) or only the Her-2/neu extracellular domain (pNeu_{ECD} and pNeu_{ECD-gDs}), respectively (see A of Fig. 1). While pNeu_{TM} or pNeu_{ECD} encodes the original Her-2/neu signal peptide sequence, the signal peptide sequence of pNeu_{TM-gDs} or
30 pNeu_{ECD-gDs} is replaced by the signal peptide sequence from glycoprotein D of herpes simplex virus type I.

Whereas injections of pNeu_{TM} or pNeu_{ECD} encoding the original signal peptide sequence induce strong Her-2/neu-specific antibody response, pNeu_{TM-gDs} or

pNeu_{ECD-gDs} encoding the signal sequence of herpes simplex virus type 1 glycoprotein D induce weak Her-2/neu-specific antibody response (see Figs. 2 and 3). However, all pNeu constructs induce similar strong Her-2/neu-specific CTL response (see Fig. 4). These constructs can be used to evaluate whether a substantial difference in the quantity of Her-2/neu-specific antibody in mice could influence protective or therapeutic immunity against Her2-CT26, a syngeneic Her-2/neu-expressing tumor.

The present invention reveals that intramuscular (i.m.) injection of pNeu_{TM}, pNeu_{ECD}, pNeu_{TM-gDs} or pNeu_{ECD-gDs} can induce complete protection against a small number of Her2-CT26 cells (see Fig. 5). Moreover, preventive antitumor efficacies of pNeu_{ECD} and pNeu_{ECD-gDs} are not significantly different even when a maximum number of tumor cells are injected subcutaneous (s.c.) or intravenous (i.v.) (see Fig. 6). This suggests strong Her-2/neu CTL response without antibody response is as effective as the collaboration of strong CTL and antibody responses in a preventive model. However, when a large number of tumor cells are preinjected in a therapeutic model, only the mouse group having both strong CTL and antibody shows a significantly improved survival rate (see Fig. 7).

The Her-2/neu expressing plasmid constructs of the present invention have the advantage of eliminating possible oncogenicity of Her-2/neu by constructing truncated Her-2/neu plasmids lacking the Her-2/neu cytoplasmic kinase domain (intracellular domain). It therefore eliminates the risks of chance transforming of normal cells and transmission of abnormal growth signal toward tumor malignancy that may be caused by tyrosine kinase in the intracellular domain. In addition, the truncated Her-2/neu of the present invention enables to avoid the dangers of autoimmunity against the Her-2/neu intracellular domain that is highly conserved among the members of the EGFR (epidermal growth factor receptor) family. It has been reported that plasmid encoding the truncated Her-2/neu are at least as effective as a plasmid encoding the total Her-2/neu (Chen, Y. et al., *Cancer Res.* 58: 1965-1971, 1998). Also, the inventive Her-2/neu expressing plasmid constructs induce the both the Her-2/neu-specific antibody response and therapeutic antitumor effect.

These results demonstrate the relative roles of CTL and antibody by DNA vaccination in a preventive model or a therapeutic model against Her-2/neu-expressing tumor. Although strong CTL activation without antibody response by DNA vaccination could achieve enough preventive efficacy against Her-2/neu-

expressing tumor challenge, DNA vaccines maximizing both arms of immune response was most beneficial in a therapeutic model.

5 To enhance the efficacy of the inventive vaccine in the clinical use, the present invention further provides Her-2/neu expressing plasmid constructs prepared by using a more efficient vector, pCK vector, in place of pTV2, to improve the expression level of Her-2/neu.

pCK vector has a stronger CMV promoter and smaller size (about 3 kb) than pTV2, and thus, a target antigen can be efficiently expressed at an increased concentration of pCK plasmid.

10 To prepare pCK plasmid constructs, the truncated Her-2/neu fragments from pNeu_{TM} and pNeu_{ECD}, which have the original Her-2/neu signal peptide and strong antitumor activity, are each inserted into pCK vector.

In another preferred embodiment of the present invention, there are provided 15 two Her-2/neu expressing plasmids based on pCK vector (pCK_{TM} and pCK_{ECD}) which encode either the Her-2/neu transmembrane and extracellular domains (pCK_{TM}) or the Her-2/neu extracellular domain only (pCK_{ECD}).

Vaccination with pCK_{TM} and pCK_{ECD} induce both strong antibody response and CTL response (see Figs. 8 and 9). The extent of immunity induced by 20 vaccination with pCK_{TM} or pCK_{ECD} is similar or slightly higher than observed for pNeu constructs. Intramuscular inoculation of pCK_{TM} and pCK_{ECD} completely prevent the growth of subcutaneous tumor and metastasis in a prevaccinated model and inhibit the tumor growth in a therapeutic model (see Figs. 10 and 11).

Her-2/neu expressing plasmid constructs of the present invention, pNeu_{TM}, 25 pNeu_{ECD}, pCK_{TM} and pCK_{ECD} have been deposited on June 26, 2002 with the Korean Culture Center of Microorganisms (KCCM) (Address: #361-221, Yurim B/D, Hongje-1-dong, Seodaemun-gu, Seoul 120-091, Republic of Korea) under the accession numbers KCCM-10393, KCCM-10394, KCCM-103395 and KCCM-10396, respectively, in accordance with the terms of Budapest Treaty on the International Recognition of the Deposit of Microorganism for the Purpose of Patent Procedure.

30 Since pCK_{TM} expressing truncated Her-2/neu is more efficient in inducing both humoral and cellular immunity, and the therapeutic antitumor activity of pCK_{TM} is slightly better than that of pCK_{ECD}, pCK_{TM} vaccination with in combination with

cytokine genes is preferred.

Accordingly, the present invention discloses the use of a cytokine as an adjuvant which is helpful for overcoming immune tolerance against Her-2/neu in tumor patients.

5 Further in purpose, the present invention chose 6 cytokines; IL-12 (Alfonso, L. C. et al., *Science* 263: 235-237, 1994), IL-15 (Min, W. et al., *Vaccine* 20: 1466-1474, 2002), IL-18 (Hanlon, L. et al., *J. Virol.* 75: 8424-8433, 2001), Eta-1, Flt3L (Mwangi, W. et al., *J. Immunol.* 169: 3837-3846, 2002), GM-CSF (Lee, A. H. et al., *Vaccine* 17: 473-479, 1999). GM-CSF and Flt3L which induce the proliferation and activation
10 of antigen presenting cells (APC) are expected to improve the delivery efficiency into APCs like dendritic cells and promote immune response including humoral and cellular immunity. IL-12, IL-15, IL-18 and Eta-1 are typical T_H1 skewing cytokines and expected to induce cell-mediated immune responses important to cancer immunity.

15 The present invention provides constructs pCK-IL12, pCK-IL15, pCK-IL18, pCK-Eta1, pCK-Flt3L and pCK-GMCSF which are obtained by inserting the respective cytokine gene into pCK vector. The effect of combining a cytokine gene adjuvant is similar to that observed for pCK_{TM} in terms of antibody production and CTL response (see Fig. 12), but coinjection of pCK_{TM} with each of the pCK-
20 cytokines, especially pCK-GMCSF, enhances the antitumor effect in the preventive and therapeutic model (see Fig. 13 and 14).

To enhance the cytokine adjuvant activity in the Her-2/neu DNA vaccination, the present invention constructed bicistronic plasmids, pCK_{TM}-GMCSF, pCK_{TM}-Flt3L, pCK_{TM}-Eta1, pCK_{TM}-IL12, pCK_{TM}-IL15, pCK_{TM}-IL18 and pCK_{TM}-IL23, in
25 which the Her-2/neu protein and each of the cytokines are translated independently. Vaccinations with the inventive bicistronic plasmids also inhibit tumor growth and metastasis (see Fig. 15 and 16). Antitumor activities of bicistronic plasmids except pCK_{TM}-IL18 are similar to those observed when two separate plasmids are coinjected. The antitumor activity of pCK_{TM}-IL18 is much higher than that of coinjection with
30 pCK_{TM} and pCK-IL18.

The above results show that the Her-2/neu expressing plasmid constructs of the present invention provide a vaccine that is not only preventive but also

therapeutic against cancers. Therefore, Her-2/neu DNA vaccines have potential usage as a therapeutic vaccine in reducing metastasis after tumor surgery or as a prophylactic vaccine for people with genetic high risk.

In accordance with another aspect of the present invention, there is also provided Her-2/neu vaccine compositions used for preventing and treating cancer.

The inventive vaccine compositions include the human Her-2/neu expressing plasmid construct of the invention and a pharmaceutically acceptable carrier. These vaccine compositions can provide protection against (used as a prophylactic) infection by the antigen induced by the human Her-2/neu expressing plasmid construct of the invention. In addition, the vaccine compositions of the invention can be used to treat (used as a therapeutic) infection by the antigen induced by the human Her-2/neu expressing plasmid construct of the invention.

The preparation of vaccine compositions that contain the human Her-2/neu expressing plasmid construct of the invention as an effective ingredient is known to one skilled in the art. Typically, such vaccines are prepared as injectables, either as liquid solutions or suspensions; solid forms suitable for solution in, or suspension in, liquid prior to infection can also be prepared. The preparation can also be emulsified, or the protein encapsulated in liposomes. The active immunogenic ingredients are often mixed with carriers which are pharmaceutically acceptable and compatible with the active ingredient. The term "pharmaceutically acceptable carrier" refers to a carrier that does not cause an allergic reaction or other untoward effect in subjects to whom it is administered. Suitable pharmaceutically acceptable carriers include, for example, one or more of water, saline, phosphate buffered saline, dextrose, glycerol, ethanol, or the like and combinations thereof. In addition, if desired, the vaccine can contain minor amounts of auxiliary substances such as wetting or emulsifying agents, pH buffering agents, and/or adjuvants which enhance the effectiveness of the vaccine. Examples of adjuvants which may be effective include but are not limited to: aluminum hydroxide, N-acetyl-muramyl-L-threonyl-D-isoglutamine (thr-MDP), N-acetyl-nor-muramyl-L-alanyl-D-isoglutamine (CGP 11637, referred to as nor-MDP), N-acetylmuramyl-L-alanyl-D-isoglutaminyl-L-alanine-2-(1'-2'-dipalmitoyl-sn-glycero-3-hydroxyphosphoryloxy)-ethylamine (CGP 19835A, referred to as MTP-PE), and RIBI, which contains three components extracted from bacteria, monophosphoryl lipid A, trehalose dimycolate and cell wall

skeleton (MPL+TDM+CWS) in a 2% squalene/Tween 80 emulsion. Other examples of adjuvants include DDA (dimethyldioctadecylammonium bromide), Freund's complete and incomplete adjuvants and QuilA. In addition, immune modulating substances such as lymphokines (e.g., IFN-g, IL-2 and IL-12) or synthetic IFN-g inducers such as poly I:C can be used in combination with adjuvants described herein.

Vaccine compositions of the present invention may be administered parenterally, by injection, for example, either subcutaneously or intramuscularly. Additional formulations which are suitable for other modes of administration include suppositories, and in some cases, oral formulations or formulations suitable for distribution as aerosols. In the case of the oral formulations, the manipulation of T-cell subsets employing adjuvants, antigen packaging, or the addition of individual cytokines to various formulation can result in improved oral vaccines with optimized immune responses. For suppositories, traditional binders and carriers may include, for example, polyalkylene glycols or triglycerides; such suppositories may be formed from mixtures containing the active ingredient in the range of 0.5 to 10%, preferably 1 to 2%. Oral formulations include such normally employed excipients as, for example, pharmaceutical grades of mannitol, lactose, starch magnesium stearate, sodium saccharine, cellulose, magnesium carbonate, and the like. These compositions take the form of solutions, suspensions, tablets, pills, capsules, sustained release formulations or powders and contain 10% to 95% of effective ingredient, preferably 25 to 70%.

The Her-2/neu expressing plasmid constructs of the present invention can be formulated into the vaccine compositions as neutral or salt forms. Pharmaceutically acceptable salts include the acid addition salts (formed with free amino groups of the peptide) and which are formed with inorganic acids such as, for example, hydrochloric or phosphoric acids, or with organic acids such as acetic, oxalic, tartaric, maleic, and the like. Salts formed with the free carboxyl groups can also be derived from inorganic bases such as, for example, sodium, potassium, ammonium, calcium, or ferric hydroxides, and such organic bases as isopropylamine, trimethylamine, 2-ethylamino ethanol, histidine, procaine, and the like.

Vaccine compositions are administered in a manner compatible with the dosage formulation, and in such amount as will be prophylactically and/or

therapeutically effective. The quantity to be administered depends on the subject to be treated, including, e.g., capacity of the subject's immune system to synthesize antibodies, and the degree of protection or treatment desired. Suitable dosage ranges are of the order of several hundred micrograms effective ingredient per vaccination with a range from about 0.01 to 10 mg/kg/day, preferably in the range from about 0.1 to 1 mg/kg/day. Suitable regimens for initial administration and booster shots are also variable but are typified by an initial administration followed by subsequent inoculations or other administrations. Precise amounts of effective ingredient required to be administered depend on the judgment of the practitioner and may be peculiar to each subject. It will be apparent to those of skill in the art that the therapeutically effective amount of Her-2/neu expressing plasmid constructs of this invention will depend, inter alia, upon the administration schedule, the unit dose of antigen administered, whether the Her-2/neu expressing plasmid construct is administered in combination with other therapeutic agents, the immune status and health of the recipient, and the therapeutic activity of the particular Her-2/neu expressing plasmid construct.

The compositions can be given in a single dose schedule, or preferably in a multiple dose schedule. A multiple dose schedule is one in which a primary course of vaccination can include 1 to 10 separate doses, followed by other doses given at subsequent time intervals required to maintain and or reinforce the immune response, for example, at 1 to 4 months for a second dose, and if needed, a subsequent dose(s) after several months. Periodic boosters at intervals of 1 to 5 years, usually 3 years, are desirable to maintain the desired levels of protective immunity.

Immunization protocols have used adjuvants to stimulate responses for many years, and as such adjuvants are well known to one of ordinary skill in the art. Some adjuvants affect the way in which antigens are presented. For example, the immune response is increased when protein antigens are precipitated by alum. Emulsification of antigens also prolongs the duration of antigen presentation.

In one aspect, an adjuvant effect is achieved by use of an agent such as alum used in about 0.05 to about 0.1% solution in phosphate buffered saline. Alternatively, the antigen is made as an admixture with synthetic polymers of sugars (Carbopol. RTM) used as an about 0.25% solution. Adjuvant effect may also be made by aggregation of the antigen in the vaccine by heat treatment with

temperatures ranging between about 70 to about 101 °C for a 30 sec to 2 min period, respectively. Aggregation by reactivating with pepsin treated (Fab) antibodies to albumin, mixture with bacterial cell(s) such as *C. parvum* or an endotoxin or a lipopolysaccharide components of Gram-negative bacteria, emulsion in physiologically acceptable oil vehicles such as mannide mono-oleate (Aracel A) or emulsion with a 20% solution of a perfluorocarbon (Fluosol-DA. RTM) used as a block substitute also may be employed.

Various polysaccharide adjuvants may also be used. For example, the use of various pneumococcal polysaccharide adjuvants on the antibody responses of mice has been described. The doses that produce optimal responses, or that otherwise do not produce suppression, should be employed as indicated. Polyamine varieties of polysaccharides are particularly preferred, such as chitin and chitosan, including deacetylated chitin.

Another adjuvant contemplated for use in the present invention is BCG. BCG (Bacillus Calmette-Guerin, an attenuated strain of *Mycobacterium*) and BCG-cell wall skeleton (CWS) may also be used as adjuvants in the invention. BCG is an important clinical tool because of its immunostimulatory properties. BCG acts to stimulate the reticulo-endothelial system, activates natural killer cells and increases proliferation of hematopoietic stem cells. Cell wall extracts of BCG have proven to have excellent immune adjuvant activity. In a typical practice of the present invention, cells of *Mycobacterium bovis* BCG are grown and harvested by methods known in the art. Besides *Mycobacterium bovis* BCG, vaccines of non-pathogenic bacteria, e. g., *Salmonella* sp., *Pseudomans* sp., *Eschericia* sp., and so on can be used in the present invention.

The present invention is further defined in the following Examples. It should be understood that these Examples, while indicating preferred embodiments of the invention, are given by way of illustration only. From the above discussion and these Examples, one skilled in the art can ascertain the essential characteristics of this invention, and without departing from the spirit and scope thereof, can make various changes and modifications of the invention to adapt it to various usage and conditions.

Reference Example 1: Cell lines and animals

The Her-2/neu expressing human breast carcinoma SK-BR3 cell line (ATCC HTB-30) and murine colon adenocarcinoma cell line CT26 (ATCC CRL-2639) were obtained from the American Type Culture Collection (Manassas, VA, USA).
5 Human breast cancer cell line SK-BR3 cells were maintained in RPMI1640 (BioWhittaker, Walkersville, MD) supplemented with 10% heat-inactivated fetal bovine serum (FBS, GIBCO, Gaithersburg, MD) and 1% penicillin-streptomycin (GIBCO). Her-2/neu-expressing transfectoma Her2-CT26 cells were prepared by transduction of CT26 cells with the cDNA-encoding human Her-2/neu (NCBI:
10 M1730). Her2/CT26 and CT26 cells were cultured in IMDM (BioWhittaker) containing 10% heat-inactivated FBS and 1% penicillin-streptomycin.

Female 5-week-old BALB/C mice were purchased from Charles River (Osaka, Japan) and kept at 22 °C, 55% relative humidity, and a daily lighting cycle of 12hrs light/ 12hrs dark with free access to food and water. The mice were housed at
15 Laboratory Animal Center of Seoul National University until use and kept in a germ-free isolator (Techniplast, Buguggiate, Italy) during the whole experiments.

Reference Example 2: Isolation of DNA plasmids for i.m. injection

20 *Escherichia coli* strain DH5 α transformed with each of the plasmids, pNeu_{TM}, pNeu_{ECD}, pNeu_{TM}-gDs, pNeu_{ECD}-gDs, pCK_{TM} and pCK_{ECD}, control vectors pTV2 and pCK, was grown in LB broth (Difco, Detroit, MI). Large-scale preparation of the plasmid DNA was carried out by the alkaline lysis method using an Endofree Qiagen Plasmid-Giga kit (Qiagen, Chatsworth, CA) according to the manufacturer's
25 instructions. DNA was then precipitated, suspended in sterile PBS (BioWhittaker) at a concentration of 2 mg/ml, and stored in aliquots at -20 °C for subsequent use in immunization protocols.

Reference Example 3: Flow cytometry (FACS)

30 To examine whether sera could specifically react Her-2/neu surface protein, SK-BR3, Her2-CT26 and CT26 cells were stripped from the culture flasks with a cell scraper (Nunc, Naperville, IL). Removed cells were washed in an FACS buffer

consisting of RPMI1640 supplemented with 2% FBS and 0.1% sodium azide. Approximately 2×10^5 cells per analysis were incubated together with a serial dilute of a serum or control antibody at 4°C for 30 min. Cells were washed 3 times a serial dilute of the same FACS buffer and then stained for 30 minutes at 4°C with an FITC-conjugated goat monoclonal antibody specific for mouse IgG (Sigma).
5 Stained cells were washed 2 times and resuspended in the same FACS buffer. To exclude dead cells from data, 1 µg/ml propidium iodide (Sigma) was added to the cell suspension and incubated for 30 sec prior to analysis. Only the cells that were negative by propidium iodide staining were gated and further analyzed for binding to
10 tumor cells. Flow cytometry was performed using a PAS IIIi flow cytometer (Partec GmbH, Münster, Germany)

Reference Example 4: Confocal microscopy for anti-Her-2/neu antibodies

15 Approximately 1×10^4 SK-BR3 cells were grown for three days on Lab-Tek chambered coverglass (Nunc, Naperville, IL) coated with 1 mg/ml poly-L-Lysine. Cells were fixed with 4% paraformaldehyde in PBS for 10 min at room temperature, washed three times with DMEM, blocked with 1% goat γ-globulin in DMEM for 1 hour at 4°C, incubated with 1:50 diluted mouse sera in a blocking solution for 8 hours
20 at 4°C, washed, and incubated with R-phycoerythrin-conjugated goat anti-mouse immunoglobulin secondary antibody (Southern Biotech, Birmingham, AL) for 30 min at room temperature. Slides were then mounted on Gel/Mount media (Fisher) and examined using a confocal microscopy (Leica TCS-SP laser scanning microscopy).

25

Reference Example 5: DNA immunization method

Briefly, each mouse received an i.m. injection of 100 µg of plasmid DNA that was dissolved in 100 µl of sterile PBS into the anterior tibialis muscle. The
30 inoculation site was pretreated with bupivacaine-HCl (ASTRA, Westborough, MA). For daily immunization for therapeutic vaccination, bupivacaine-HCl was pretreated only once just before the first immunization. Sera were collected via the retro-orbital plexus at selected time points and monitored for the presence of anti-Her-

2/neu antibodies.

Reference Example 6: Chromium-release assays

5 Splenocytes prepared by extracting spleen from immunized mice were cultured with mytomycin-C treated Her2-CT26 cells for 6 days, and were assayed for the lysis of CT26 or Her2-CT26 target cells in a 4 hour ^{51}Cr -release assay.

Her2-CT26 or CT26 tumor target cells were labeled with ^{51}Cr by incubating 2×10^6 cells with 200 μCi $\text{Na}^{51}\text{CrO}_4$ (NEN Research Products, Boston, MA) in 200 μl saline at 37°C for 90 min. The unincorporated ^{51}Cr was removed by four washes with RPMI1640. Graded numbers of effector cells were mixed with 10000 labeled target cells in 200 μl RPMI plus 10% FBS in the wells of a round-bottom microtiter plate. The plate was incubated at 37°C for 4 hours. After the incubation, the plate was centrifugated, and a 100 μl aliquot was removed from each well for counting with a γ -scintillation counter (Packard, Minaxi Auto Gamma 5000 Series). The percent lysis was calculated by formula 1:

<Formula 1>

$$\text{percent specific lysis (\%)} = 100 \times [(\text{cpm}_{\text{experimental}} - \text{cpm}_{\text{spontaneous}}) / (\text{cpm}_{\text{max}} - \text{cpm}_{\text{spontaneous}})]$$

The cpm_{max} value was determined by adding 10 μl of 5% triton-X (Sigma) to wells containing ^{51}Cr -labeled target cells. Each group contained a duplicate. The $\text{cpm}_{\text{spontaneous}}$ value was determined by adding only an equal volume of the medium without the addition of splenocytes or triton-X.

Reference Example 7: Tumor challenge

Mice were challenged by injection with Her2-CT26 cells suspended in sterile PBS either subcutaneously on the flank or intravenously. The three-dimensional size of each tumors was measured with a caliper, and the volume was calculated by formula 2:

<Formula 2>

$$\text{tumor volume (mm}^3\text{)} = (\text{width} \times \text{length} \times \text{depth}) \text{ mm}^3 \times \pi/6$$

Animals were monitored twice a week for the development of palpable tumors. Mice showing any symptom of acute sickness, hard to breathe or rare movement were sacrificed.

Example 1: Construction of Her-2/neu expressing plasmids

pTV2 and pTV2-gDs (Lee, S. W. et al., *J. Virol.* 72:8430-8436, 1998) and pCK (Lee Y., et. al., *Biochem Biophys Res Commun.* 272:230-235, 2000; Deposit Accession No: KCCM-10179) were used as expression vectors. pTV2-gDs is an expression vector which was cloned to contain the signal sequence of herpes simplex virus type 1 glycoprotein D in expression vector pTV2. The cDNA encoding the entire human Her-2/neu gene (SEQ ID NO: 1) was inserted into the pRC/CMV backbone (Invitrogen, San Diego, CA) to produce a full-length Her-2/neu plasmid (9.6 Kb).

The plasmid pNeu_{ECD}, encoding the extracellular domain of Her-2/neu without the intracellular and transmembrane domains of Her-2/neu, was generated from the PCR product of the full-length Her-2/neu plasmid using NF6 (SEQ ID NO: 4) and NSR1 (SEQ ID NO: 5) as a primer pair, and cloned into the KpnI and XbaI sites of pTV2. Similarly, the plasmid pNeu_{TM}, encoding the extracellular and transmembrane domains of Her-2/neu, was generated from the PCR product of the full-length Her-2/neu plasmid using NF5 (SEQ ID NO: 6) and NRM2 (SEQ ID NO: 7) as a primer pair, and cloned into the KpnI and XbaI sites of pTV2 (Fig. 1).

The plasmid pNeu_{ECD-gDs}, encoding the extracellular domain of Her-2/neu without the intracellular and transmembrane domains of Her-2/neu, was generated from the PCR product of the full-length Her-2/neu plasmid using NSF2 (SEQ ID NO: 8) and NSR1 (SEQ ID NO: 5) as a primer pair, and cloned into the AscI and XbaI sites of pTV2-gDs. Similarly, the plasmid pNeu_{TM-gDs}, encoding the extracellular and transmembrane domains of Her-2/neu, was generated from the PCR product of the full-length Her-2/neu plasmid using NF3 (SEQ ID NO: 9) and NRM2 (SEQ ID NO: 7) as a primer pair, and cloned into the AscI and XbaI sites of pTV2-gDs. The

plasmids pCK_{ECD} and pCK_{TM} were prepared by inserting into the KpnI-XbaI site of pCK vector truncated Her-2/neu gene fragments obtained from pNeu_{ECD} and pNeu_{TM}, respectively. PCR was carried out at 94°C for 2 min; 94°C for 15 sec, 55°C for 30 sec and 68°C for 3.5 min; and 72°C for 7 min.

Thus generated were four Her-2/neu expressing plasmids (pNeu_{TM}, pNeu_{ECD}, pNeu_{TM-gDs}, and pNeu_{ECD-gDs}), each encoding both the Her-2/neu transmembrane and extracellular domains (pNeu_{TM} and pNeu_{TM-gDs}) or only the Her-2/neu extracellular domain (pNeu_{ECD} and pNeu_{ECD-gDs}) (Fig 1a). While pNeu_{TM} or pNeu_{ECD} encoded the original Her-2/neu signal peptide sequence, the signal peptide sequence of pNeu_{TM-gDs} or pNeu_{ECD-gDs} was replaced by the signal peptide sequence from glycoprotein D of herpes simplex virus type I.

Example 2: Induction of anti-Her-2/neu Antibody by pNeu constructs vaccination

Tests were conducted to examine whether various pNeu plasmid constructs could induce anti-Her-2/neu antibodies as follows.

Each mouse prepared in Reference Example 1 received three i.m. injections of 100 µg of plasmid DNA prepared in Reference Example 2 according to a preset immunization schedule (Fig. 1b). Some mice of each group were sacrificed and the lytic function of Her-2/neu-specific CTL was determined. Other mice were challenged with Her-2/neu expressing tumor for evaluating antitumor immunity. Sera were obtained from BALB/c mice before the first injection and one week after the third vaccination, and the anti-Her-2/neu antibody titer in the serum was measured based on its binding to the breast cancer cell line, SK-BR3, using a flow cytometry. Her-2/neu-specific serum IgG titers of all mice vaccinated with pNeu_{TM}, pNeu_{ECD}, pNeu_{TM-gDs}, or pNeu_{ECD-gDs} were determined and presented based on the greatest dilution of serum for which a shift in the mean fluorescence intensity of binding affinity to SK-BR3 cells was seen relative to an irrelevant control antibody.

<Table 1>

pTV2 (n=5)	pNeu _{TM} (n=5)	pNeu _{ECD} (n=5)	pNeu _{TM-gDs} (n=5)	pNeu _{ECD-gDs} (n=5)
<50	12800	12800	800	<50

<50	12800	12800	50	<50
<50	3200	12800	<50	<50
<50	12800	12800	800	<50
<50	3200	12800	50	<50

As shown in Table 1, the observed IgG titers were ranked in the order of $p\text{Neu}_{\text{ECD}} > p\text{Neu}_{\text{TM}} > p\text{Neu}_{\text{TM-gDs}} > p\text{Neu}_{\text{ECD-gDs}} = p\text{TV2}$. As expected, none of the sera collected from animals before the injection of plasmid DNA had detectable anti-Her-2/neu binding activities. Moreover, none of the animals injected with pTV2 made detectable anti-Her-2/neu antibodies at 1:50 dilution. However, vaccination with pNeu_{TM} or pNeu_{ECD} resulted in high Her-2/neu-specific IgG titers (Fig. 2, A) and serum samples diluted by 1:800 produced a wide shift in the mean fluorescence intensity (Fig. 2, B and C). In contrast, vaccination with pNeu_{TM-gDs} or pNeu_{ECD-gDs} resulted in a low or undetectable IgG titer and serum samples diluted by 1:50 revealed a little or a barely detectable shift in the mean fluorescence intensity (Fig. 2, D and E).

The existence of Her-2/neu-specific antibodies in mouse sera immunized with pNeu_{TM} or pNeu_{ECD-gDs} was also confirmed by confocal microscopic analysis. Mouse serum immunized with pNeu_{TM} (Fig. 3, B) demonstrated clear localization of anti-Her-2/neu antibodies on the surface of SK-BR3 that was not shown with the mouse sera immunized with pTV2 (Fig. 3, A) or pNeu_{ECD-gDs} (Fig. 3, C), which is consistent with the anti-Her-2/neu antibody titers presented in Fig. 2.

Example 3: Induction of Her-2/neu-specific CTL by pNeu constructs vaccination

Having demonstrated that vaccination with pNeu constructs boosted high to very low Her-2/neu-specific antibody responses in vaccinated mice (Fig. 2), Her-2/neu-specific CTL responses induced in the same mice were evaluated as follows.

Splenocytes were prepared 2 weeks after the third immunization from the same mice that were tested for Her-2/neu-specific serum IgG titers. Splenocytes were cultured with mytomycin-C-treated human Her-2/neu expressing syngeneic murine transfectoma, Her2-CT26 cells for 6 days, and were assayed for the lysis of CT26 or Her2-CT26 target cells by a 4-h ⁵¹Cr-release assay.

As a result, splenocytes from mice vaccinated with pNeu_{TM} (Fig. 4, B), pNeu_{ECD} (Fig. 4, C), pNeu_{TM-gDs} (Fig. 4, D), or pNeu_{ECD-gDs} (Fig. 4, E) exhibited CTL-dependent lysis of Her2-CT26 that was not shown with splenocytes from pTV2 vaccinated control mice (Fig. 4, A) and the relative strength of Her-2/neu-specific CTL response was in order of pNeu_{TM} > pNeu_{ECD} > pNeu_{TM-gDs} > pNeu_{ECD-gDs} > pTV2. Percent Her-2/neu-specific lysis by splenocytes from mice immunized with any one of pNeu constructs was comparable to the others of pNeu constructs and were 80 ~ 90% at an E:T ratio of 50:1 and 60 ~ 70% at an E:T ratio of 10:1 (Fig. 3, B to E). However, splenocytes from any group of mice did not induce CTL-dependent lysis of CT26 cells.

In brief, all Her-2/neu expressing plasmids induced strong Her-2/neu-specific CTL response, which was irrelevant to their signal peptide sequences. However, they induced substantially different Her-2/neu-specific antibody responses according to their signal peptide sequences. Only pNeu_{TM} and pNeu_{ECD} with the original signal sequence showed high Her-2/neu-specific IgG titers (Fig. 2). When their signal sequence was replaced by a viral signal sequence, pNeu_{TM-gDs} generated a low level of anti-Her-2/neu antibodies, and pNeu_{ECD-gDs}, a very low level of anti-Her-2/neu antibodies.

Example 4: Prevention of tumor growth by pNeu constructs vaccination

Antitumor immunity against human Her-2/neu expressing syngeneic murine tumor cell line Her2-CT26 in mice was evaluated as follows.

Initially, titration studies were performed to determine the optimal number of tumor cells to be injected s.c. or i.v. into mice to generate subcutaneous tumor formation or lung metastasis, and the results showed that Her2-CT26 cells induced subcutaneous or lung metastatic tumor in BALB/c mice when 5×10^4 cells or more were injected s.c. or i.v.. Since a long survival period may help to distinguish antitumor efficacy of Her-2/neu DNA plasmids, 5×10^4 cells were chosen as the initial cell number for i.v. or s.c. tumor challenge. Each mouse received three i.m. injections of 100 μ g plasmid DNA according to a preset immunization schedule (Fig. 1b) and one and a half weeks after the third injection of plasmid DNA, each mouse was challenged i.v. or s.c. with 5×10^4 Her2-CT26 cells.

In the above subcutaneous tumor model study, all of the animals injected with pTV2 developed palpable tumors (Fig. 5, A). On the other hand, tumors were completely suppressed in all groups of mice each injected with pNeu_{TM}, pNeu_{ECD}, pNeu_{TM-gDs}, or pNeu_{ECD-gDs} for 60 days following s.c. tumor injection. In a metastasis model, all group of mice injected with either pNeu_{TM}, pNeu_{ECD}, pNeu_{TM-gDs}, or pNeu_{ECD-gDs} survived i.v. tumor challenge (Fig. 5, B). However, four of the seven mice (57%) injected with pTV2 and all mice injected with only PBS did not survive lung metastasis.

Example 5: Comparison of antitumor immunity by pNeu_{ECD} and pNeu_{ECD-gDs}

Examples 2 to 4 demonstrated contrasting differences of Her-2/neu-specific antibody titers but comparable CTL responses in mice immunized with different pNeu plasmids. In addition, all groups of mice each immunized with pNeu_{TM}, pNeu_{ECD}, pNeu_{TM-gDs}, or pNeu_{ECD-gDs} rejected 5×10^4 s.c. tumor challenge. Since the number of tumor cells that were injected s.c. or i.v. into mice was too small to induce tumor in immunized mice, it was very difficult to distinguish antitumor efficacies by the difference in immune response induced by different pNeu constructs. Therefore, the number of tumor cells to be injected was increased by a factor of 100 (5×10^6) for s.c. tumor challenge and by a factor of 40 (2×10^6) for i.v. tumor challenge relative to that of tumor cells in the first tumor experiment to evaluate the relative importance of Her-2/neu-specific antibody and CTL toward inhibition of Her2-CT26. It was impossible to use a cell number of more than 2×10^6 for i.v. tumor challenge because there was the danger of blood vessel blockage by excessive tumor cells injected i.v. Chosen for a comparative purpose was a set of pNeu_{ECD} and pNeu_{ECD-gDs} that generated the largest difference in Her-2/neu-specific antibody titers among the four different Her-2/neu-expressing plasmids. Each mouse received three i.m. injections of 100 μ g plasmid DNA according to the same immunization schedule (Fig. 1, B), and 10 days after the third injection of plasmid DNA, each mouse was challenged s.c. with 5×10^6 or i.v. with 2×10^6 Her2-CT26.

In the subcutaneous model, all eight animals injected with pTV2 developed tumors and the mean tumor volume reached over 2000 mm³ before day 19 post s.c. tumor challenge (Fig. 6, A). The mean tumor volume of eight mice injected with

pNeu_{ECD} was 82.2 mm³ at day 23 and that of eight mice injected with pNeu_{ECD-gDs} was 67.9 mm³. While there was significant suppression of tumor growth in mice injected with pNeu_{ECD} ($p = 2.9900 \times 10^{-8}$, Student's t test) or pNeu_{ECD-gDs} ($p = 2.8400 \times 10^{-8}$, Student's t test), the difference in the mean tumor volume between the two immunized groups was not statistical significance ($P = 0.8684$, Student's t test). In the metastasis model, lung metastasis was inhibited until day 40 in eight of the eight mice (100%) injected with pNeu_{ECD} and in seven of the eight mice (88%) injected with pNeu_{ECD-gDs} (Fig. 6, B). All mice vaccinated with pTV2 did not survived lung metastasis. Again, although the survival was significantly prolonged by treatment with pNeu_{ECD} ($p < 0.0001$, Mantel-Haenszel test) or pNeu_{ECD-gDs} ($p = 0.0002$, Mantel-Haenszel test) compared with pTV2, there was no significant difference between pNeu_{ECD} and pNeu_{ECD-gDs} ($p = 0.3173$, Mantel-Haenszel test).

Example 6: Efficacy of pNeu constructs vaccination in a therapeutic model

Preventive model tumor experiments were performed by challenging immunized mice with tumor cells. To compare the antitumor efficacies of pNeu_{ECD} and pNeu_{ECD-gDs} in a therapeutic model, mice were challenged with tumor cells first, and then received i.m. injections of DNA plasmids. 6-week old naive mice were challenged i.v. with 1×10^5 or 5×10^5 Her2-CT26 cells, and then were divided into 4 groups. Beginning 1 hour after the tumor injection, each mouse received the first i.m. injection of 100 μ g of pNeu_{ECD} or pNeu_{ECD-gDs}, followed by four more daily i.m injections with the same DNA plasmid.

The results in Figs. 5 and 6 show that when 1×10^5 tumor cells were injected, all mice treated with pNeu_{ECD} or pNeu_{ECD-gDs} survived lung metastasis for the following 40 days (Fig. 7, A). However, five of the eight mice (63%) injected with pTV2 and all mice (100%) injected with only PBS did not survive lung metastasis. Although pNeu_{ECD} and pNeu_{ECD-gDs} improved the survival rate significantly ($p = 0.0085$, Mantel-Haenszel test) as compared with pTV2, there was no significant difference between pNeu_{ECD} and pNeu_{ECD-gDs}.

On the other hand, when the number of tumor cells was increased 5 times (5×10^5), only the mice injected with pNeu_{ECD} exhibited an increased survival rate which was statistically significantly ($p = 0.0237$, Mantel-Haenszel test) compared

with mice injected with pTV2 (Fig. 7, B). However, the mice injected with pNeu_{ECD-gDs} did not show significantly enhanced survival ($p = 0.4628$, Mantel-Haenszel test) as compared with the mice injected with pTV2. Nonetheless, consistently with the preventive model, there was no significant difference in antitumor immunity between pNeu_{ECD} and pNeu_{ECD-gDs} ($p = 0.4263$, Mantel-Haenszel test).

In summary, therapeutic efficacies of Her-2/neu DNA vaccines were evaluated by changing the number of preinjected tumor cells. When mice were treated with a small number of metastatic tumor cells, both pNeu_{ECD} and pNeu_{ECD-gDs} prolonged the survival period significantly and there was no significant difference between their antitumor immunity. However, when a large number of tumor cells were used, only pNeu_{ECD} improved the survival rate.

Example 7: Comparative analysis of immune response induced by pNeu constructs and pCK constructs

To enhance the clinical efficacy of vaccine, Her-2/neu DNA plasmid vector was constructed with pCK vector which has stronger promoter activity than pTV2. The KpnI-XbaI fragments of truncated Her-2/neu genes obtained from pNeu_{ECD} and pNeu_{TM} were each inserted into the KpnI-XbaI site of pCK vector. Thus, pCK_{TM} expressing the extracellular and transmembrane domains and pCK_{ECD} expressing the extracellular domain of Her-2/neu were prepared.

To evaluate the immunogenicity of pCK_{TM} and pCK_{ECD}, BALB/c mice were vaccinated with pCK_{TM}, pCK_{ECD}, pNeu_{ECD} and pNeu_{TM} and the sera and spleen were obtained from immunized mice 10 days after the third intramuscular inoculation with each DNA plasmid. SK-BR3 cells were incubated with 400 fold-diluted sera, followed by binding with FITC conjugated goat anti-mouse IgG. Estimation of Her-2/neu specific antibody response was performed by end-point titration using a flow cytometer. The result in Fig. 8 shows that vaccination of mice with pCK_{TM} and pCK_{ECD} induced Her-2/neu-specific IgG antibody responses similar to vaccination with pNeu constructs, wherein uncolored and colored histograms represent control antibody and diluted serum, respectively.

Furthermore, Her-2/neu-specific CTL activity was assayed against Her2-CT26

in a standard ^{51}Cr -release assay. Vaccination of mice with pCK_{TM} and pCK_{ECD} also induce strong CTL responses (Fig. 9). CTL responses induced by vaccination of pCK constructs were slightly higher than when vaccinated with pNeu constructs.

5 **Example 8: Antitumor activity of pCK_{TM} and pCK_{ECD}**

To determine the antitumor effect of pCK constructs of Her-2/neu, female BALB/c mice were vaccinated intramuscularly three times with 100 μg PBS, pCK, pCK_{ECD} or pCK_{TM} in two-week intervals, respectively. The mice were challenged s.c. or i.v. with 1×10^6 Her2-CT26 2 weeks after final vaccination. The three-dimensional size of grown solid tumor induced by s.c. injection of Her2-CT26 was measured with a caliper. The number of live mice was counted everyday and the results were presented as the percentage of live mice per treatment group.

10 Growth of solid tumors induced by s.c. injection of Her2-CT26 was inhibited completely in the mice vaccinated with pCK_{TM} or pCK_{ECD} (Fig. 10, A). In the lung metastasis model, pCK_{TM} and pCK_{ECD} prolonged the survival period, demonstrating strong suppression of lung metastasis (Fig. 10, B). In summary, protective immunity against Her2-CT26 challenge also could be achieved by vaccination with pCK_{TM} or pCK_{ECD}.

20 To test therapeutic effects of pCK_{TM} and pCK_{ECD}, mice were vaccinated intramuscularly with 100 μg PBS, pCK, pCK_{ECD} or pCK_{TM}, 1 hr after 2×10^5 Her2-CT26 i.v. challenge. The number of live mice was counted everyday and the results were presented as the percentage of live mouse per treatment group. Post-vaccination with pCK_{TM} or pCK_{ECD} was effective for the protection against the growth of metastatic colony and appeared to inhibit death by lung metastasis (Fig. 11). Thus, preventive antitumor effects of pNeu constructs were retained in pCK_{TM} and pCK_{ECD}. Furthermore, in the therapeutic model to evaluate protective immunity against preinjected tumor cells, pCK_{TM} and pCK_{ECD} prolonged the survival rate significantly. Since therapeutic antitumor activity of pCK_{TM} was slightly better than that of pCK_{ECD}, pCK_{TM} was chosen as a model for Her-2/neu DNA vaccine in combination with cytokine genes.

30 **Example 9: Immune responses and antitumor activities induced by coinjection of**

pCK_{TM} and various cytokine plasmids

To use cytokine genes as a molecular adjuvant in the Her-2/neu DNA vaccination, six cytokine gene-contained pCK vectors, pCK-GMCSF, pCK-IL12, pCK-IL15, pCK-IL18, pCK-Eta1 and pCK-Flt3L, were prepared as follows. GM-CSF and Flt3L, which promote the proliferation and activation of antigen presenting cells, are expected to improve the delivery efficiency into professional antigen presenting cells like dendritic cells and to increase immune responses. IL-12, IL-15, IL-18 and Eta-1 are representative T_H1 skewing cytokines and expected to induce cell-mediated immune responses important to cancer immunity.

Eta-1 (SEQ ID NO: 10), IL-18 (SEQ ID NO: 11), IL-15 (SEQ ID NO: 12) and Flt3L (SEQ ID NO: 13) genes were amplified from mRNA isolated from the spleen of BALB/c mice by RT-PCR (SUPERScriptTM II RT, GIBCO BRL) with specific primers (Eta-1, with EF1 of SEQ ID NO: 14 and ER1 of SEQ ID NO: 15; IL-18, with 18F1 of SEQ ID NO: 16 and 18R1 of SEQ ID NO: 17; IL-15, with 15F1 of SEQ ID NO: 18 and 15R1 of SEQ ID NO: 19; and Flt3L, with FF1 of SEQ ID NO: 20 and FR1 of SEQ ID NO: 21) according to the manufacturer's instructions. Cloned cytokine genes were inserted into pCK to generate pCK-Eta1, pCK-IL18, pCK-IL15 and pCK-Flt3L. pCK-GMCSF and pCK-IL12 were constructed by inserting the EcoRI-XbaI and XhoI fragments of pTV2-GMCSF (Cho, J. H. et al., *Vaccine* 17: 1136-1144, 1999) and pTV2-IL12 (Ha, S. J. et al., *Nat. Biotechnol.* 20: 381-386, 2002), into pCK vector, respectively.

To analyze the effects of cytokine gene adjuvants in the antibody production and CTL response, mice were intramuscularly injected with pCK_{TM} and each of the pCK-cytokines (Fig. 12a). 3 weeks after the final vaccination, antibody titration was performed by a flow cytometry to determine Her-2/neu-specific antibody production and ⁵¹Cr-release assay, to measure CTL responses.

<Table 2>

Mice	Her-2/neu specific IgG titer							
	PCK	pCK _{TM}	+IL-12	+IL-15	+IL-18	+Eta-1	+Flt3L	+GM-CSF
1	<50	3200	6400	6400	6400	3200	800	3200
2	<50	3200	3200	3200	6400	3200	3200	6400

3	<50	6400	6400	1600	6400	6400	800	6400
4	<50	3200	400	1600	12800	6400	1600	1600

As shown in Table 2, Her-2/neu-specific antibodies were sufficiently produced by vaccination of pCK_{TM} with or without cytokines but no significant difference was found for the groups coinjected with cytokine gene plasmids (Fig. 12b).

5

<Table 3>

	Effector : Target ratio		
	3:1	10:1	30:1
pCK	5.5±0.69	8.8±3.16	20.3±5.43
pCK _{TM}	52.6±6.03	67.6±0.56	76.4±1.21
pCK _{TM} + pCK-IL12	34.6±5.67	59.3±8.82	79.3±6.99
pCK _{TM} + pCK-IL15	39.0±0.76	53.6±1.08	70.7±6.30
pCK _{TM} + pCK-IL18	21.8±1.44	38.8±1.53	47.7±2.55
pCK _{TM} + pCK-Eta1	59.5±9.01	88.7±11.07	96.2±4.52
pCK _{TM} + pCK-Flt3L	48.4±2.99	79.6±3.22	95.9±2.38
pCK _{TM} + pCK-GMCSF	34.6±12.96	50.6±15.56	64.3±13.8

Table 3 shows a summary of the CTL responses observed in Fig. 12b. As shown in Table 3, the percentage of target lysis increased slightly by vaccination of pCK_{TM} with pCK-Eta1 or pCK-Flt3L but decreased slightly by vaccination of pCK_{TM} with pCK-IL18 or pCK-GMCSF. Nonspecific lysis using CT26 as target cell was not found in all mice.

10

Example 10: Antitumor activity induced by coinjection of pCK_{TM} and cytokine plasmids

15

To determine antitumor activity induced by coinjection of pCK_{TM} and a cytokine plasmid, preventive and therapeutic experiments were performed using BALB/c mice. As shown in Fig. 13a and Fig. 14a, BALB/c mice were challenged with Her2-CT26 cells after and before vaccination with pCK_{TM} and each of the cytokine

20

plasmids. BALB/c mice were co-injected i.m. with 100 μ g pCK_{TM} and 100 μ g each of the pCK-cytokine plasmids. The mice were challenged i.v. or s.c. with 1×10^6 Her2-CT26 at week 3 after the 2nd vaccination. Tumor growth was measured with a caliper twice a week and the volume was calculated for each mouse.

5 The growth of subcutaneous tumor was inhibited by covaccination with pCK_{TM} and a cytokine plasmid, especially pCK-GMCSF, pCK-Eta1 and pCK-IL15 (Fig. 13b). Metastases of intravenously challenged Her2-CT26 were inhibited by vaccination with pCK_{TM} and pCK-GMCSF (Fig. 13c and 13d).

10 Mice were vaccinated intramuscularly with 100 μ g pCK_{TM} and each of the pCK-cytokine plasmids after 2×10^5 Her2-CT26 i.v. challenge. The number of live mice was counted everyday and the results were presented as the percentage of live mouse per treatment group. Co-vaccination with pCK_{TM} and pCK-cytokine plasmids except pCK-Eta1 improved the survival rate more than when vaccinated only with pCK_{TM} (Fig. 14b and 14c).

15 Therefore, the preventive antitumor activity of pCK_{TM} was promoted by co-injection of a particular cytokine plasmid such as pCK-GMCSF both in tumor growth model and metastasis model.

20 Example 11: Construction of bicistronic plasmids expressing Her-2/neu and cytokine

To enhance the antitumor activity of Her-2/neu DNA vaccination, constructed were bicistronic plasmids, pCK_{TM}-GMCSF, pCK_{TM}-Flt3L, pCK_{TM}-Eta1, pCK_{TM}-IL12, pCK_{TM}-IL15, pCK_{TM}-IL18 and pCK_{TM}-IL23, in which the Her-2/neu protein and
25 each of the cytokines had been translated independently. The internal ribosomal entry site (IRES) of encephalomyocarditis virus (EMCV) between the Her-2/neu gene and cytokine gene enabled the simultaneous expression of Her-2/neu protein and cytokine (Fig. 15a).

30 To generate bicistronic plasmids co-expressing Her-2/neu and cytokine proteins, GM-CSF, Flt3L, IL-15, IL-18 and Eta-1 genes were amplified by PCR using specific primers as described in Example 9, which were inserted downstream of IRES of EMCV of pCK_{TM}-IRES. IRES of EMCV having the nucleotide sequence of SEQ ID NO: 22 was derived from pCK-IL12. For IL-12 and IL-23 (Belladonna, M. L., et

al., *J. Immunol.* 168: 5448-5454, 2002), IRES was amplified by PCR using pCK-IL12 as a template and IRES-F1 of SEQ ID NO: 23 and IRES-R1 of SEQ ID NO: 24 as a primer pair, and the amplified product was inserted into the NotI-XhoI site of pCK_{TM} to obtain pCK_{TM}-IRES.

5

Example 12: Antitumor effects induced by bicistronic plasmids expressing Her-2/neu and cytokine

10 To evaluate preventive antitumor activities of pCK_{TM}-cytokine plasmids, mice were vaccinated with each of the seven pCK_{TM}-cytokine plasmids (pCK_{TM}-GMCSF, pCK_{TM}-Flt3L, pCK_{TM}-Eta1, pCK_{TM}-IL12, pCK_{TM}-IL15, pCK_{TM}-IL18 and pCK_{TM}-IL23) according to the vaccination schedule shown in Fig. 15b.

15 Intramuscular inoculation with pCK_{TM}-cytokine constructs inhibited more thoroughly the growth of tumor implanted subcutaneously than when only pCK was injected (Fig. 15c). Especially, pCK_{TM}-IL18, pCK_{TM}-GMCSF, pCK_{TM}-IL12 and pCK_{TM}-Flt3L exhibited outstanding effects in the inhibition of tumor growth. In the tumor metastasis model, pCK_{TM}-IL18 completely protected mice from lung metastasis (Fig. 15d). Anti-metastatic effects of other bicistronic constructs were similar to the case of pCK_{TM}. Protection against metastasis of preinjected Her2-
20 CT26 was assayed in the treatment model according to the schedule shown in Fig. 14a. Vaccination with pCK_{TM}-GMCSF and pCK_{TM}-IL18 prolonged the survival rate (Fig. 16, A and B). Collectively, pCK_{TM}-GMCSF and pCK_{TM}-IL18 appeared to markedly increase tumor suppression effects in the preventive and therapeutic model.

25

While the embodiments of the subject invention have been described and illustrated, it is obvious that various changes and modifications can be made therein without departing from the spirit of the present invention which should be limited only by the scope of the appended claims.

BUDAPEST TREATY ON THE INTERNATIONAL
RECOGNITION OF THE DEPOSIT OF MICROORGANISMS
FOR THE PURPOSES OF PATENT PROCEDURE

INTERNATIONAL FORM

To: Chang-Yuik Kang
PanGenomics, Inc., Innovation Center,
Institute of Molecular Biology and Genetics,
Seoul National University, Sillimdong,
Kwanakgu, Seoul 151-057, Korea

RECEIPT IN THE CASE OF AN ORIGINAL
issued pursuant to Rule 7.1 by the
INTERNATIONAL DEPOSITARY AUTHORITY
identified at the bottom of this page

I. IDENTIFICATION OF THE MICROORGANISM	
Identification reference given by the DEPOSITOR : pNeuTM	Accession number given by the INTERNATIONAL DEPOSITARY AUTHORITY: KCCM-10393
II. SCIENTIFIC DESCRIPTION AND/OR PROPOSED TAXONOMIC DESIGNATION	
The microorganism identified under I above was accompanied by:	
<input type="checkbox"/> a scientific description <input type="checkbox"/> a proposed taxonomic designation (Mark with a cross where applicable)	
III. RECEIPT AND ACCEPTANCE	
This International Depositary Authority accepts the microorganism identified under I above, which was received by it on Jun. 26. 2002. (date of the original deposit) ¹	
IV. INTERNATIONAL DEPOSITARY AUTHORITY	
Name : Korean Culture Center of Microorganisms Address : 361-221, Yurim B/D Hongje-1-dong, Seodaemun-gu SEOUL 120-091 Republic of Korea	Signature(s) of person(s) having the power to represent the International Depositary Authority or of authorized official(s): Date: Jul. 3. 2002

¹ Where Rule 6.4(d) applies, such date is the date on which the status of international depositary authority was acquired : where a deposit made outside the Budapest Treaty after the acquisition of the status of international depositary authority is converted into a deposit under the Budapest Treaty, such date is the date on which the microorganism was received by the international depositary authority.


Sole page

BUDAPEST TREATY ON THE INTERNATIONAL
RECOGNITION OF THE DEPOSIT OF MICROORGANISMS
FOR THE PURPOSES OF PATENT PROCEDURE

INTERNATIONAL FORM

To: Chang-Yuil Kang
PanGenomics, Inc., Innovation Center,
Institute of Molecular Biology and Genetics,
Seoul National University, Sillimdong,
Kwanakgu, Seoul 151-057, Korea

RECEIPT IN THE CASE OF AN ORIGINAL
issued pursuant to Rule 7.1 by the
INTERNATIONAL DEPOSITARY AUTHORITY
identified at the bottom of this page

I. IDENTIFICATION OF THE MICROORGANISM	
Identification reference given by the DEPOSITOR : pNeuECD	Accession number given by the INTERNATIONAL DEPOSITARY AUTHORITY: KCCM-10394
II. SCIENTIFIC DESCRIPTION AND/OR PROPOSED TAXONOMIC DESIGNATION	
The microorganism identified under I above was accompanied by: <input type="checkbox"/> a scientific description <input type="checkbox"/> a proposed taxonomic designation (Mark with a cross where applicable)	
III. RECEIPT AND ACCEPTANCE	
This International Depositary Authority accepts the microorganism identified under I above, which was received by it on Jun. 26. 2002. (date of the original deposit) ¹	
IV. INTERNATIONAL DEPOSITARY AUTHORITY	
Name : Korean Culture Center of Microorganisms Address : 361-221, Yurim B/D Hongje-1-dong, Seodae-mun-gu SEOUL 120-091 Republic of Korea	Signature(s) of person(s) having the power to represent the International Depositary Authority or of authorized official(s): Date: Jul. 3. 2002 <div style="text-align: center; margin-top: 10px;">  </div>


¹ Where Rule 6.4(d) applies, such date is the date on which the status of international depositary authority was acquired : where a deposit made outside the Budapest Treaty after the acquisition of the status of international depositary authority is converted into a deposit under the Budapest Treaty, such date is the date on which the microorganism was received by the international depositary authority.

BUDAPEST TREATY ON THE INTERNATIONAL
RECOGNITION OF THE DEPOSIT OF MICROORGANISMS
FOR THE PURPOSES OF PATENT PROCEDURE

INTERNATIONAL FORM

To. Chang-Yuil Kang
PanGenomics, Inc., Innovation Center,
Institute of Molecular Biology and Genetics,
Seoul National University, Sillimdong,
Kwanakgu, Seoul 151-057, Korea

RECEIPT IN THE CASE OF AN ORIGINAL
issued pursuant to Rule 7.1 by the
INTERNATIONAL DEPOSITARY AUTHORITY
identified at the bottom of this page

I. IDENTIFICATION OF THE MICROORGANISM	
Identification reference given by the DEPOSITOR : pCK-ECD	Accession number given by the INTERNATIONAL DEPOSITARY AUTHORITY: KCCM-10395
II. SCIENTIFIC DESCRIPTION AND/OR PROPOSED TAXONOMIC DESIGNATION	
The microorganism identified under I above was accompanied by: <input type="checkbox"/> a scientific description <input type="checkbox"/> a proposed taxonomic designation (Mark with a cross where applicable)	
III. RECEIPT AND ACCEPTANCE	
This International Depositary Authority accepts the microorganism identified under I above, which was received by it on Jun. 26. 2002. (date of the original deposit) ¹	
IV. INTERNATIONAL DEPOSITARY AUTHORITY	
Name : Korean Culture Center of Microorganisms Address : 361-221, Yurim B/D Hongje-1-dong, Seodaemun-gu SEOUL 120-091 Republic of Korea	Signature(s) of person(s) having the power to represent the International Depositary Authority or of authorized official(s): Date: Jul. 3. 2002 <div style="text-align: right;">  </div>

¹ Where Rule 6.4(d) applies, such date is the date on which the status of international depositary authority was acquired : where a deposit made outside the Budapest Treaty after the acquisition of the status of international depositary authority is converted into a deposit under the Budapest Treaty, such date is the date on which the microorganism was received by the international depositary authority.

Sole page

BUDAPEST TREATY ON THE INTERNATIONAL
RECOGNITION OF THE DEPOSIT OF MICROORGANISMS
FOR THE PURPOSES OF PATENT PROCEDURE

INTERNATIONAL FORM

To: Chang-Yuil Kang
PanGenomics, Inc., Innovation Center,
Institute of Molecular Biology and Genetics,
Seoul National University, Sillimdong,
Kwanakgu, Seoul 151-057, Korea

RECEIPT IN THE CASE OF AN ORIGINAL
issued pursuant to Rule 7.1 by the
INTERNATIONAL DEPOSITARY AUTHORITY
identified at the bottom of this page

I. IDENTIFICATION OF THE MICROORGANISM	
Identification reference given by the DEPOSITOR : pCK-TM	Accession number given by the INTERNATIONAL DEPOSITARY AUTHORITY: KCCM-10396
II. SCIENTIFIC DESCRIPTION AND/OR PROPOSED TAXONOMIC DESIGNATION	
The microorganism identified under I above was accompanied by: <input type="checkbox"/> a scientific description <input type="checkbox"/> a proposed taxonomic designation (Mark with a cross where applicable)	
III. RECEIPT AND ACCEPTANCE	
This International Depositary Authority accepts the microorganism identified under I above, which was received by it on Jun. 26. 2002. (date of the original deposit) ¹	
IV. INTERNATIONAL DEPOSITARY AUTHORITY	
Name : Korean Culture Center of Microorganisms Address : 361-221, Yurim B/D Hongje-1-dong, Seodaemun-gu SEOUL 120-091 Republic of Korea	Signature(s) of person(s) having the power to represent the International Depositary Authority or of authorized official(s): Date: Jul. 3. 2002

¹ Where Rule 6.4(d) applies, such date is the date on which the status of international depositary authority was acquired : where a deposit made outside the Budapest Treaty after the acquisition of the status of international depositary authority is converted into a deposit under the Budapest Treaty, such date is the date on which the microorganism was received by the international depositary authority.

What is claimed is:

1. An Her-2/neu plasmid construct having anti-cancer activity which is prepared by inserting a truncated human Her-2/neu gene lacking the intracellular domain into plasmid pTV2 or pCK.
5
2. The plasmid construct of claim 1, wherein the human Her-2/neu gene has the nucleotide sequence of SEQ ID NO: 2.
- 10 3. The plasmid construct of claim 2, which is pNeu_{TM} (KCCM-10393) or pCK_{TM} (KCCM-10396)
4. The plasmid construct of claim 1, wherein the truncated human Her-2/neu gene further lacks the transmembrane domain.
- 15 5. The plasmid construct of claim 4, wherein the human Her-2/neu gene has the nucleotide of SEQ ID NO: 3.
6. The plasmid construct of claim 5, which is pNeu_{ECD} (KCCM-10394) or
20 pCK_{ECD} (KCCM-10395).
7. The plasmid construct of claim 1, wherein the signal peptide of the human Her-2/neu gene is replaced by the signal peptide of herpes simplex type I glycoprotein D (gD).
- 25 8. The plasmid construct of claim 7, which is pNeu_{TM-gDs}.
9. The plasmid construct of claim 4, wherein the signal peptide of the human Her-2/neu gene is replaced by the signal peptide of herpes simplex type I
30 glycoprotein D (gD).
10. The plasmid construct of claim 7, which is pNeu_{ECD-gDs}.

11. The plasmid construct of claim 1, which further translates a cytokine gene besides the human Her-2/neu gene.
12. The plasmid construct of claim 11, wherein the cytokine gene is selected from the group consisting of granulocyte-macrophage colony-stimulating factor (GM-CSF), FMS-like tyrosine kinase 3 ligand (Flt3L), early T lymphocyte activation-1 (Eta-1), interleukin-12 (IL-12), IL-15 and IL-18.
13. A DNA vaccine for preventing and/or treating cancer, which comprises the plasmid construct of claim 1 as an effective ingredient and a pharmaceutically acceptable carrier.
14. The DNA vaccine of claim 13, which further comprises a cytokine gene expressing plasmid.
15. The DNA vaccine of claim 14, wherein the cytokine gene is selected from the group consisting of GM-CSF, Flt3L, Eta-1, IL-12, IL-15 and IL-18.
16. A method for preventing and/or treating cancer, which comprises the step of administering an effective amount of the DNA vaccine of claim 13.

1/18
Fig. 1a





Fig. 1b



2/18
Fig. 2










Fig. 3




3/18
Fig. 4


4/18
Fig. 5


5/18
Fig. 6

6/18
Fig. 7

7/18
Fig. 8



8/18
Fig. 9

9/17
Fig. 10




10/18
Fig. 11

Fig. 12a




11/18
Fig. 12b

Her2-CT26



CT26





12/18
Fig. 13a

Fig. 13b





13/18
Fig. 13c

Fig. 13d





14/18
Fig. 14a

Fig. 14b








15/18
Fig. 14c

Fig. 15a



17/18
Fig. 15d

18/18
Fig. 16

SEQUENCE LISTINGS

<110> PANGENOMICS Co., Ltd

<120> Her-2/neu DNA VACCINE HAVING ANTI-CANCER ACTIVITY

<130> PCA30540/PAN/PCT

<150> KR2002-41764

<151> 2002-07-16

<150> KR2003-38012

<151> 2003-06-12

<160> 24

<170> KopatentIn 1.71

<210> 1

<211> 4530

<212> DNA

<213> human Her-2/neu gene cDNA

<400> 1
aattctcgag ctctcgacc ggtcgacgag ctcgagggtc gacgagctcg gggcgcgcg 60
cccgcccccc acccctcgca gcaccccgcg ccccgcgccc tcccagcggg gtccagccgg 120
agccatgggg ccggagccgc agtgagcacc atggagctgg cggccttggt cggctggggg 180
ctctcctcg ccctcttgcc ccccgagacc gcgagcacc aagtgtgcac cggcacagac 240
atgaagctgc ggctccctgc cagtcccgag acccacctgg acatgctccg ccacctctac 300
cagggtgccc aggtggtgca gggaaacctg gaactcacct acctgcccac caatgccagg 360
ctgtccttcc tgcaggatat ccaggagggtg cagggtacg tgctcatcgc tcacaaccaa 420
gtgaggcagg tccactgca gaggctgagg attgtgcgag gcaccagct cttgaggac 480
aactatgccc tggccgtgct agacaatgga gaccgctga acaataccac ccctgtcaca 540
ggggcctccc caggaggcct gcgggagctg cagcttcgaa gcctcacaga gatcttgaaa 600
ggaggggtct tgatccagcg gaacccccag ctctgctacc aggacacgat tttgtggaag 660

gacatcttcc acaagaacaa ccagctggct ctcacactga tagacaccaa ccgctctcgg 720
gcctgccacc cctgttctcc gatgtgtaag ggctcccgct gctggggaga gattctgag 780
gattgtcaga gcctgacgcg cactgtctgt gccggiggct gtgcccgtg caaggggcca 840
ctgcccactg actgtgcca tgagcagtgt gctgccggct gcacggggcc caagcactct 900
gactgcctgg cctgcctcca cttaaccac agtggcatct gtgagctgca ctgcccagcc 960
ctggtcacct acaacacaga cacgtttgag tccatgcca atcccgaggg ccggtataca 1020
ttcggcgcca gctgtgtgac tgctgtccc tacaactacc tttctacgga cgtgggatcc 1080
tgcaccctcg tctgccccct gcacaaccaa gaggtgacag cagaggatgg aacacagcgg 1140
tgtgagaagt gcagcaagcc ctgtgccga gtgtgtatg gtctgggcat ggagcacttg 1200
cgagaggatga gggcagttac cagtccaat atccaggagt ttgtggctg caagaagatc 1260
tttgggagcc tggcatittt gccggagagc ttgatgggg acccagcctc caacactgcc 1320
ccgctccagc cagagcagct ccaagtgttt gagactctgg aagagatcac aggttaccta 1380
tacatctcag catggccgga cagcctgcct gacctcagcg tcttcagaa cctgcaagta 1440
atccggggac gaattctgca caatggcgcc tactcgctga cctgcaagg gctgggcac 1500
agctggctgg ggcctgcgtc actgaggga ctgggcagt gactggccct catccaccat 1560
aacaccacc tctgttctgt gcacacggtg ccctgggacc agctctttcg gaaccgcac 1620
caagctctgc tccacactgc caaccggcca gaggacgagt gtgtggcga gggcctggcc 1680
tgccaccagc tgtgcgcccg agggcactgc tgggtccag ggcccacca gtgtgtcaac 1740
tgcagccagt tccitcgggg ccaggagtgc gtggaggaat gccagtgact gcaggggctc 1800
cccaggagat atgtgaatgc caggcactgt ttgccgtgcc accctgagtg tcagcccag 1860
aatggctcag tgacctgttt tggaccggag gctgaccagt gtgtggcctg tgcccactat 1920
aaggaccctc cttctgcgt ggcccgtgc ccagcgggtg tgaaacctga cctctcctac 1980
atgcccactct ggaagtittc agatgaggag ggcgatgcc agccttgccc catcaactgc 2040

accactcct gtgtggacct ggatgacaag ggctgccccg ccgagcagag agccagccct 2100
ctgacgtcca tcgtctctgc ggtgggtggc attctgctgg tcgtggtctt gggggtggtc 2160
tttgggatcc tcatcaagcg acggcagcag aagatccgga agtacacgat gcggagactg 2220
ctgcaggaaa cggagctggt ggagccgctg acacctagcg gagcgatgcc caaccaggcg 2280
cagatgcgga tcctgaaaga gacggagctg aggaaggtga aggtgcttgg atctggcgct 2340
tttggcacag tctacaaggg catctggatc cctgatgggg agaattgtgaa aattccagtg 2400
gccatcaaag tgttgagggg aaacacatcc cccaaagcca acaaagaaat cttagacgaa 2460
gcatacgtga tggctgggtg gggctcccca tatgtctccc gccttctggg catctgcctg 2520
acatccacgg tgcagctggt gacacagctt atgccctatg gctgcctctt agaccatgtc 2580
cgggaaaacc gcggacgcct gggctcccag gacctgctga actggtgtat gcagattgcc 2640
aaggggatga gctacctgga ggatgtgcgg ctogtacaca gggacttggc cgctcggaac 2700
gtgctggtca agagtcccaa ccatgtcaaa attacagact tcgggctggc tcggctgctg 2760
gacattgacg agacagagta ccatgcagat gggggcaagg tgcccatcaa gtggatggcg 2820
ctggagtcca ttctccgccg gcggttcacc caccagagtg atgtgtggag ttatggtgtg 2880
actgtgtggg agctgatgac ttttggggcc aaaccttacg atgggatccc agcccgggag 2940
atccctgacc tgctggaaaa gggggagcgg ctgcccagc ccccatctg caccattgat 3000
gtctacatga tcatggtcaa atgttggatg attgactctg aatgtcggcc aagattccgg 3060
gagttggtgt ctgaattctc cgcgatggcc agggacccc agcgcttctt ggtcatccag 3120
aatgaggact tgggcccagc cagtcccttg gacagcacct tctaccgctc actgctggag 3180
gacgatgaca tgggggacct ggtggatgct gaggagtatc tggtaacca gcagggttc 3240
ttctgtocag acctgcccc gggcgctggg ggcatggtcc accacaggca ccgcagctca 3300
tctaccagga gtggcggtgg ggacctgaca ctagggtgg agccctctga agaggaggcc 3360
cccaggctc cactggcacc ctccgaaggg gctggctccg atgtattga tggtagctg 3420

ggaatggggg cagccaaggg gctgcaaagc ctccccacac atgaccccag ccctctacag 3480
 cggtagctg aggaccccac agtaccctg ccctctgaga ctgatggcta cgttgcccc 3540
 ctgacctgca gccccagcc tgaatatgtg aaccagccag atgttcggcc ccagccccct 3600
 tcgccccgag agggccctct gcctgctgcc cgacctgtg gtgccactct ggaaagggcc 3660
 aagactctct cccagggaa gaatggggc gtcaaagacg ttttgcctt tgggggtgcc 3720
 gtggagaacc ccgagtactt gacaccccag ggaggagctg cccctcagcc ccacctct 3780
 cctgccttca gccagcctt cgacaacctc tttactggg accaggacc accagagcgg 3840
 ggggctccac ccagcacctt caaagggaca cctacggcag agaaccaga gtacctgggt 3900
 ctggacgtgc cagtgtgaac cagaaggcca agtccgcaga agccctgatg tctctcagg 3960
 gagcaggaa ggcctgactt ctgctggcat caagaggtag gagggccctc cgaccactc 4020
 caggggaacc tgccatgcca ggaacctgtc ctaaggaacc ttcttctctg cttagattcc 4080
 cagatggctg gaaggggtcc agcctcgttg gaagaggaac agcactgggg agtctttgtg 4140
 gattctgagg cctgcccac tgagactcta ggtccagtg gatccacag cccagcttgg 4200
 ccctttcctt ccagatcctg ggtactgaaa gccttaggga agctggcctg agaggggaag 4260
 cggccctaag ggagtgtcta agaacaaaag cgaccattc agagactgtc cctgaaacct 4320
 agtactgcc cccatgagga aggaacagca atggtgtcag tatccaggct ttgtacagag 4380
 tgctttctg tttagttttt actttttttg tttgttttt ttaaagacga aataaagacc 4440
 caggggagaa tgggtgttgt atggggaggg aagtgtgggg ggtccttctc cacaccact 4500
 ttgtccattt gcaaataat tttggaaaac 4530

<210> 2
 <211> 2052
 <212> DNA
 <213> human Her-2/neu gene without intracellular region

<400> 2
 atggagctgg cggccttgtg ccgctggggg ctctcctcg ccctcttggc cccggagacc 60

gcgagcacc aagtgtgcac cggcacagac atgaagctgc ggctccctgc cagtcccag 120
accacactgg acatgtccg ccacctctac cagggctgcc aggtggtgca gggaaacctg 180
gaactcacct acctgcccac caatgccagc ctgtcttcc tgcaggatat ccaggaggtg 240
cagggctacg tgcctatgc tcacaaccaa gtgaggcagg tccactgca gaggtgcgg 300
attgtgcgag gcaccagct ctttgaggac aactatgcc tggcgtgct agacaatgga 360
gacccgtga acaataccac cctgtcaca ggggcctccc caggaggcct gcgggagctg 420
cagcttcgaa gcctcacaga gatcttgaaa ggaggggtct tgaiccagcg gaacccccag 480
ctctgtacc aggacacgat tttgtggaag gacatctcc acaagaaca ccagctggct 540
ctcactga tagacacaa ccgtctcgg gcctgccacc cctgttctcc gatgtgtaag 600
ggctccgct gctggggaga gattctgag gattgtcaga gcctgacg cgactgtctgt 660
gccggtggct gtgccgctg caaggggcca ctgccactg actgtgcc tgcagctgt 720
gctgccggct gcacggggcc caagcactct gactgcctgg cctgcctcca ctcaaccac 780
agtggcatct gtgagctgca ctgccagcc ctggctacct acaacacaga cacgtttgag 840
tccatgcca atcccgagg cgggtatata ttccgggcca gctgtgtgac tgcctgtccc 900
tacaactacc tttctacgga cgtgggatcc tgcacctcg tctgccccct gcacaacaa 960
gaggtgacag cagaggatgg aacacagcg tgtgagaagt gcagcaagcc ctgtgccga 1020
gtgtgctatg gtctgggcat ggagcacttg cgagaggtga gggcagttac cagtccaat 1080
atccaggagt ttgctggctg caagaagatc tttgggagcc tggcatttct gccggagagc 1140
tttgatggg acccagcctc caacactgcc ccgtccagc cagagcagct ccaagtgtt 1200
gagactctgg aagagatcac aggtacctt tacatctcag catggccgga cagcctgcct 1260
gacctcagcg tctccagaa cctgcaagta atccggggac gaattctgca caatggcgcc 1320
tactcgtga ccctgcaagg gctgggcatc agctggctgg ggctgcgctc actgaggaa 1380
ctgggcagtg gactggccct catccacat aacaccacc tctgttctg gcacacggtg 1440

ccctgggacc agctctttcg gaaccogcac caagctctgc tccacactgc caaccggcca 1500
 gaggacgagt gtgtggcgga gggcctggcc tgccaccagc tgtgcgcccg agggcactgc 1560
 tgggttccag ggcccaccca gtgtgtcaac tgcagccagt tccttcgggg ccaggagtgc 1620
 gtggaggaat gccgagtact gcaggggctc cccagggagt atgtgaatgc caggcactgt 1680
 ttgccgtgcc accctgagtg tcagccccag aatggctcag tgacctgttt tggaccggag 1740
 gctgaccagt gtgtggcctg tgcccactat aaggaccctc ccttctgcgt ggcccgtgc 1800
 cccagcggtg tgaacctga cctctctac atgcccactt ggaagtctcc agatgaggag 1860
 ggcgcattgcc agccttgccc catcaactgc acccactcct gtgtggacct ggatgacaag 1920
 ggctgccccg ccgagcagag agccagccct ctgacgtcca tcgtctctgc ggtgggtggc 1980
 attctgctgg tcgtggcttt gggggtggtc ttggggatcc tcatcaagcg acggcagcag 2040
 aagatccgga ag 2052

<210> 3
 <211> 1956
 <212> DNA
 <213> human Her-2/neu gene without intracellular region and transmembrane domain

<400> 3
 atggagctgg cggccttgct cgcctggggg ctctctctcg ccctcttgcc ccccgagacc 60
 gccagacccc aagtgtgcac cggcacagac atgaagctgc ggctccctgc cagtcccgag 120
 acccacctgg acatgctccg ccacctctac cagggtgcc aggtgggtgca gggaaacctg 180
 gaactcacct acctgcccac caatgccagc ctgtccttcc tgcaggatat ccaggagggtg 240
 cagggtacg tgctcatgc tcacaaccaa gtgaggcagg tcccactgca gaggctgcgg 300
 attgtgcgag gcaccagct cttgaggac aactatgcc tggccgtgct agacaatgga 360
 gaccgctga acaataccac ccctgtcaca ggggcctccc caggaggcct gcgggagctg 420
 cagcttcgaa gcctcacaga gatcttgaaa ggaggggtct tgatccagcg gaacccccag 480

ctctgctacc aggacacgat ttttggaag gacatcttcc acaagaacaa ccagctggct 540
ctcacactga tagacaccaa ccgctctcgg gcctgccacc cctgttctcc gatgtgtaag 600
ggctcccgt gctggggaga gatttctgag gattgtcaga gcctgacgcg cactgtctgt 660
gccggtggct gtgcccgtg caaggggcca ctgccactg actgctgccca tgagcagtgt 720
gttgccggct gcacggggcc caagcactct gactgcctgg cctgcctcca cttaaccac 780
agtggcatct gtgagctgca ctgccagcc ctggtcacct acaacacaga cacgtttgag 840
tccatgcca atcccaggg ccgtataca ttccggcca gctgtgtgac tgcctgtccc 900
tacaactacc tttctacgga cgtgggatcc tgcacctcg tctgccccct gcacaacaa 960
gagtgacag cagaggatgg aacacagcgg tgtgagaagt gcagcaagcc ctgtgccga 1020
gtgtgctatg gtctgggcat ggagcacttg cgagaggatga gggcagttac cagtgccaat 1080
atccaggagt ttgctggctg caagaagatc ttggggagcc tggcatttct gccggagagc 1140
tttgatgggg acccagcctc caacactgcc ccgtccagc cagagcagct ccaagtgttt 1200
gagactctgg aagagatcac aggttaccta tacatctcag catggccgga cagcctgcct 1260
gacctcagcg tcttcagaa cctgcaagta atccggggac gaattctgca caatggcgcc 1320
tactcgctga cctgcaagg gctgggcata agctggctgg ggctgcgctc actgaggaa 1380
ctgggcagt gactggccct catccacat aacaccacc tctgcttctg gcacacggtg 1440
ccctgggacc agctctttcg gaaccgcac caagctctgc tccacactgc caaccggcca 1500
gaggacgagt gtgtgggca gggcctggcc tgccaccagc tgtgcgcccg agggcactgc 1560
tgggtccag ggcccacca gtgtgtcaac tgcagccagt tccttcgggg ccaggagtgc 1620
gtggaggaat gccgagtact gcaggggctc ccaggaggat atgtgaatgc caggcactgt 1680
ttgccgtgcc accctgagtg tcagccccag aatggctcag tgacctgttt tggaccggag 1740
gctgaccagt gtgtggcctg tgccactat aaggaccctc ccttctgcgt ggcccgtgc 1800
cccagcggtg tgaacctga cctctctac atgcccactt ggaagtctcc agatgaggag 1860

ggcgcatgcc agccttgccc catcaactgc accactcct gtgtggacct ggatgacaag 1920

ggctgccccg ccgagcagag agccagccct ctgacg 1956

<210> 4
<211> 27
<212> DNA
<213> Artificial Sequence

<220>
<223> NF6 primer

<400> 4 27
ggtacatgg agctggcggc cttgtgc

<210> 5
<211> 31
<212> DNA
<213> Artificial Sequence

<220>
<223> NSR1 primer

<400> 5 31
gtctagatga ttcacgtcag agggctggct c

<210> 6
<211> 23
<212> DNA
<213> Artificial Sequence

<220>
<223> NF5 primer

<400> 6 23
gcagtggtag ccaagcttag cac

<210> 7

<211> 27
<212> DNA
<213> Artificial Sequence

<220>
<223> NRM2 primer

<400> 7
ttctagagca gtctccgcat cgtctac

27

<210> 8
<211> 28
<212> DNA
<213> Artificial Sequence

<220>
<223> NSF2 primer

<400> 8
ggcgcgcccc ggcacagaca tgaagctg

28

<210> 9
<211> 24
<212> DNA
<213> Artificial Sequence

<220>
<223> NF3 primer

<400> 9
gccgcagcgg ccgccatgga gctg

24

<210> 10
<211> 1535
<212> DNA
<213> mouse Eta-1 gene

<400> 10
gggggggggg gggggggggg ggggctttcc ttgctcctta tgagaggtgg agaggtagaa

60

aaggcacaca aatattgact cactgaaatt ttctctgaga ttagaaaaga ttccataaat 120
tattggtgac ttggtggtga tctagtgtg ccaagaggt gtttgaacct gacaagacat 180
caactgtgcc tcataaaata tgttgcagga ctaactacga ccatgagatt ggcagtatt 240
tgcttttgcc tgtttggcat tgcttctcc ctcccggtga aagtactga ttctggcagc 300
tcagaggaga agctttacag cctgcacca gatcctatag ccacatggcc ggtgcctgac 360
ccatctcaga agcagaatct ccttgcgcca cagaatgtg tgtcctctga agaaaaggat 420
gactttaagc aagaaactct tccaagcaat tccaatgaaa gccatgacca catggacgac 480
gatgatgacg atgatgatga cgatggagac catgcagaga gcgaggattc tgtggactcg 540
gatgaatctg acgaatctca ccattcggat gagtctgatg agaccgtcac tgctagtaca 600
caagcagaca ctttactcc aatcgtccct acagtcatg tcccaacgg ccgaggtgat 660
agcttggctt atggactgag gtcaaagtct aggagtctcc aggtttctga tgaacagtat 720
cctgatgcca cagatgagga cctcacctct cacatgaaga gcggtgagtc taaggagtcc 780
ctcgaatgca tccctgttgc ccagcttctg agcatgccct ctgatcagga caacaacgga 840
aagggcagcc atgagtcaag tcagctggat gaaccaagtc tggaacaca cagacttgag 900
cattccaaag agagccagga gagtgcgat cagtcggatg tgatcgaatg tcaagcaagt 960
tccaaagcca gcctggaaca tcagagccac aagtctcaca gccacaagga caagctagtc 1020
ctagacccta agagtaagga agatgatagg tatctgaaat tccgaatttc tcatgaatta 1080
gagagttcat ctctgaggt caactaaaga agaggcaaaa acacagttcc ttactttgca 1140
tttagtaaaa acaagaaaaa gtgttagtga ggttaagca ggaatactaa ctgctcattt 1200
ctcagttcag tggatatatg tatgtagaga aagagaggta atattttggg ctcttagctt 1260
agtctgttgt ttcatgcaa caccgttgta accaaaagct tctgcacttt gcttctgtg 1320
ttcctgtaca agaaatgcaa cggccactgc attttaatga ttgtattct ttcatgaata 1380
aatgtatgt agaaataagt aaatttactg aaacaagcaa gaattaaaag agaaactgta 1440

acagtctata tcactataacc cttttagttt tataatttagc atatattttg ttgtgattta 1500
tttttttttg ttggtgtgaa taaatcttgt taacg 1535

<210> 11
<211> 1535
<212> DNA
<213> mouse IL-18 gene

<400> 11
gggggggggg gggggggggg ggggctttcc ttgctcctta tgagaggtagg agaggtagaa 60
aaggcacaca aatattgact cactgaaatt ttctctgaga ttagaaaga ttccataaat 120
tattggtgac ttggtggtga tctagtggg ccaagagtgt gtttgaacct gacaagacat 180
caactgtgcc tcataaaata tgttgcagga ctaactacga ccatgagatt ggcagtgatt 240
tgcttttgcc tgtttggcat tgccctctcc ctcccggtag aagtgactga ttctggcagc 300
tcagaggaga agctttacag cctgcaccca gatcctatag ccacatggcc ggtgcctgac 360
ccatctcaga agcagaatct ccttgggcca cagaatgctg tgcctctga agaaaaggat 420
gactttaagc aagaaactct tccaagcaat tccaatgaaa gccatgacca catggacgac 480
gatgatgacg atgatgatga cgatggagac catgcagaga gcgaggattc tgtggactcg 540
gatgaatctg acgaatctca ccattcggat gagtctgatg agaccgtcac tgctagtaca 600
caagcagaca ctttcaactcc aatcgtccct acagtcatg tcccaacgg ccgaggatg 660
agcttggctt atggactgag gtcaaagtct aggagtttcc aggtttctga tgaacagtat 720
cctgatgcca cagatgagga cctcacctct cacatgaaga gcggtgagtc taaggagtcc 780
ctcgatgtca tccctgttgc ccagcttctg agcatgccct ctgatcagga caacaacgga 840
aagggcagcc atgagtcaag tcagctggat gaaccaagtc tggaaacaca cagacttgag 900
cattccaaag agagccagga gagtgccgat cagtcggatg tgatcgatag tcaagcaagt 960
tccaaagcca gcctggaaca tcagagccac aagtttcaca gccacaagga caagctagtc 1020
ctagacccta agagtaagga agatgatagg tatctgaaat tccgaatttc tcatgaatta 1080

gagagttcat ctctgaggt caactaaaga agaggcaaaa acacagttcc ttactttgca 1140
 tttagtaaaa acaagaaaaa gtgttagtga gggtaagca ggaatactaa ctgctcattt 1200
 ctgagttcag tggatataatg tatgtagaga aagagaggta atattttggg ctcttagctt 1260
 agtctgttgt ttcattgcaa caccgttgta accaaaagct tctgcacttt gcttctgttg 1320
 ttctgtaca agaaatgcaa cgccactgc attttaatga ttgttattct ttcattgaata 1380
 aaatgtatgt agaaataagt aaatttactg aaacaagcaa gaattaaaag agaaactgta 1440
 acagtctata tcaactatacc cttttagttt tataattagc atatattttg ttgtgattta 1500
 ttttttttg ttggtgtgaa taaatcttgt taacg 1535

<210> 12
 <211> 1250
 <212> DNA
 <213> mouse IL-15

<400> 12
 ctctgtcca gccactcttc ccagagttc tcttcttcat cctccccctt gcagagtagg 60
 gcagcttgca ggtctctctg caagtctctc ccaattctct gcgccccaaa gacttgcagt 120
 gcatctcctt acgcgctgca gggaccttgc cagggcagga ctgccccgc ccagttgcag 180
 agttggacga agacgggata ctgctgtgtt tggaaggctg agttccacat ctaacagctc 240
 agagaggta ggaagaatac caccttgaca catggccctc tggctcttca aagcactgcc 300
 tcttcatggt ccttgcctgt gaggtcctta agaacacaga aacctatgtc agcagataac 360
 cagcctacag gaggccaaga agagttctgg atggatggca gctggaagcc catcgccata 420
 gccagctcat cttaacatt gaagctctta cctgggcatt aagtaatgaa aattttgaaa 480
 ccataatga ggaatacatc catctcgtgc tacttgtgtt tccttctaaa cagtcacttt 540
 ttaactgagg ctggcattca tgtcttcatt ttgggctgtg tcagtgtagg tctccctaaa 600
 acagaggcca actggataga tgtaagatat gacctggaga aaattgaaag ccttattcaa 660

tctattcata ttgacaccac ttatacact gacagtgact ttcattcccag ttgcaaagtt 720
 actgcaatga actgctttct cctggaattg caggttattt tacatgagta cagtaacatg 780
 actcttaatg aaacagtaag aaacgtgctc taccttgcaa acagcactct gtcttctaac 840
 aagaatgtag cagaatctgg ctgcaaggaa tgtgaggagc tggaggagaa aaccttcaca 900
 gagtttttgc aaagctttat acgcattgtc caaatgttca tcaacacgtc ctgactgcat 960
 gcgagcctct tccgtgtttc tgttattaag gtacctccac ctgctgctca gaggcagcac 1020
 agctccatgc atttgaaatc tgctgggcaa actaagcttc ctaacaagga gataatgagc 1080
 cacttggatc acatgaaatc ttggaaatga agagaggaaa agagctcgtc tcagacttat 1140
 ttttgcctgc ttatttttaa ttatttgctt catttgtaca tatttgtaat ataacagaag 1200
 atgtggaata aagttgtatg gatattttat caattgaaat ttaaaaaaaaa 1250

<210> 13
 <211> 699
 <212> DNA
 <213> mouse Flt3L gene

<400> 13
 atgacagtgc tggcgccagc ctggagccca aattcctccc tgttgctgct gttgctgctg 60
 ctgagtcctt gcctgcgggg gacacctgac tgttacttca gccacagtcc catctcctcc 120
 aacttcaaag tgaagttag agagttgact gaccacctgc ttaaagatta cccagtcact 180
 gtggccgtca atcttcagga cgagaagcac tgcaaggcct tgtggagcct ctctcctagcc 240
 cagcgctgga tagagcaact gaagactgtg gcagggctta agatgcaaac gcttctggag 300
 gacgtcaaca ccgagataca ttttgtcacc tcatgtacct tccagcccct accagaatgt 360
 ctgcgattcg tccagaccaa catctccac ctctgaagg acacctgcac acagctgctt 420
 gctctgaagc cctgtatcgg gaaggcctgc cagaatttct ctcggtgcct ggaggtgcag 480
 tgccagccgg actcctccac cctgtgccc ccaaggagtc ccatagccct agaagccacg 540
 gagctcccag agcctcggcc caggcagctg ttgctcctgc tgctgctgct gctgcctctc 600

acactgggtgc tgctggcagc cgcctggggc cttcgctggc aaagggcaag aaggaggggg 660
gagctccacc ctgggggtgcc cctcccctcc catccctag 699

<210> 14
<211> 21
<212> DNA
<213> Artificial Sequence

<220>
<223> EF1 primer

<400> 14 21
ctggtaccat gagattggca g

<210> 15
<211> 22
<212> DNA
<213> Artificial Sequence

<220>
<223> ER1 primer

<400> 15 22
cctctagatt agttgacctc ag

<210> 16
<211> 25
<212> DNA
<213> Artificial Sequence

<220>
<223> 18F1 primer

<400> 16 25
tgaattcatg gctgccatgt cagaa

<210> 17

<211> 23
<212> DNA
<213> Artificial Sequence

<220>
<223> 18R1 primer

<400> 17
ttctagacta actttgatgt aag 23

<210> 18
<211> 28
<212> DNA
<213> Artificial Sequence

<220>
<223> 15F1 primer

<400> 18
tgaattcatg aaaattttga aaccatat 28

<210> 19
<211> 35
<212> DNA
<213> Artificial Sequence

<220>
<223> 15R1 primer

<400> 19
ttctagacta aaagctttgc aaaaactctg tgaag 35

<210> 20
<211> 24
<212> DNA
<213> Artificial Sequence

<220>
<223> FF1 primer

<400> 20
tgaattcatg acagtctgg cgcc

24

<210> 21
<211> 24
<212> DNA
<213> Artificial Sequence

<220>
<223> FR1 primer

<400> 21
ttctagacta ctgcctgggc cgag

24

<210> 22
<211> 600
<212> DNA
<213> IRES sequence from pCK-mIL12

<400> 22
ggatccgata agcttgatat cgaattccgc ccccccccc cctaacgtta ctggccgaag 60
ccgcttggaa taaggccggt gtgcgtttgt ctatatgtta tttccacca tattgccgtc 120
ttttggcaat gtgagggccc ggaaacctgg ccctgtcttc ttgacgagca ttcctagggg 180
tctttccct ctcgccaag gaatgcaagg tctgttgaat gtcgtgaagg aagcagttcc 240
tctggaagct tcttgaagac aaacaacgtc tgtagcgacc ctttcagggc agcggaaccc 300
cccacctggc gacaggtgcc tctgcggcca aaagccacgt gtataagata cacctgcaaa 360
ggcggcaciaa cccagtgcc acgttgtgag ttggatagtt gtggaaagag tcaaattggct 420
ctcctcaagc gtattcaaca aggggctgaa ggatgccaa aaggtacccc attgtatggg 480
atctgatctg gggcctcggg gcacatgctt tacatgtgtt tagtccaggt taaaaaacg 540
tctaggcccc ccgaaccacg gggacgtggt tttcctttga aaaacacgat aataccatgg 600

<210> 23

<211> 31
<212> DNA
<213> Artificial Sequence

<220>
<223> IRES-F1 primer

<400> 23
gcggccgcga taagcttgat atogaattcc g

31

<210> 24
<211> 27
<212> DNA
<213> Artificial Sequence

<220>
<223> IRES-R1 primer

<400> 24
ctcgagtatt atcgtgtttt tcaaagg

27

INTERNATIONAL SEARCH REPORT

International application No.
PCT/KR03/01400

A. CLASSIFICATION OF SUBJECT MATTER**IPC7 C12N 15/85**

According to International Patent Classification (IPC) or to both national classification and IPC

B. FIELDS SEARCHED

Minimum documentation searched (classification system followed by classification symbols)

IPC7 C12N 15/85

Documentation searched other than minimum documentation to the extent that such documents are included in the fields searched
Korean patents and applications for inventions since 1975

Electronic data base consulted during the international search (name of data base and, where practicable, search terms used)
NCBI pubmed database, NCBI blastn database, Delphion Research Intellectual Property network database

C. DOCUMENTS CONSIDERED TO BE RELEVANT

Category*	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
A	Urbanelli L. et al 'Targeted gene transduction of mammalian cells expressing the HER2/neu receptor by filamentous phage' In Journal of Molecular Biology, 2001, vol.313, No.5, p965-76 (NCBI accession No:NM_004448) see the whole document	1,2,3-12,13-15,16
A	Coussens L. et al 'Tyrosine kinase receptor with extensive homology to EGF receptor shares chromosomal location with neu oncogene' In Science. 1985, vol.230, No.4730, p1132-9 (NCBI accession No:M11730) see the whole document	1,2,3-12,13-15,16
A	Yamamoto T. et al 'Similarity of protein encoded by the human c-erb-B-2 gene to epidermal growth factor receptor' In Nature. 1986, vol.319, No.6050, p230-4. (NCBI accession No:X03363) see the whole document	1,2,3-12,13-15,16

☐ Further documents are listed in the continuation of Box C.

☐ See patent family annex.

* Special categories of cited documents:

"A" document defining the general state of the art which is not considered to be of particular relevance

"E" earlier application or patent but published on or after the international filing date

"L" document which may throw doubts on priority claim(s) or which is cited to establish the publication date of citation or other special reason (as specified)

"O" document referring to an oral disclosure, use, exhibition or other means

"P" document published prior to the international filing date but later than the priority date claimed

"T" later document published after the international filing date or priority date and not in conflict with the application but cited to understand the principle or theory underlying the invention

"X" document of particular relevance; the claimed invention cannot be considered novel or cannot be considered to involve an inventive step when the document is taken alone

"Y" document of particular relevance; the claimed invention cannot be considered to involve an inventive step when the document is combined with one or more other such documents, such combination being obvious to a person skilled in the art

"&" document member of the same patent family

Date of the actual completion of the international search

28 NOVEMBER 2003 (28.11.2003)

Date of mailing of the international search report

28 NOVEMBER 2003 (28.11.2003)

Name and mailing address of the ISA/KR



Korean Intellectual Property Office
920 Dunsan-dong, Seo-gu, Daejeon 302-701,
Republic of Korea

Facsimile No. 82-42-472-7140

Authorized officer

LIM, Hea Joon

Telephone No. 82-42-481-5590

